

# Analysis on groups

Problem Set 3

8 March 2018

## Exercise 1

Let  $X$  be a set and  $m \in \ell^\infty(X)^*$ . Consider the following properties:

- (i)  $m(f) \geq 0$  if  $f \geq 0$ .
- (ii)  $m(\mathbf{1}_X) = 1$ , where  $\mathbf{1}_X$  is the constant function 1.
- (iii)  $\|m\| = 1$ .

Show that each of these properties follows from the two other.

## Exercise 2

The goal of this exercise is to prove that the group  $\mathrm{SL}_2(\mathbf{Z})$  contains a free group  $F_2$ .

Consider the elements  $u = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$  and  $v = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$ . Analyse again what they do to a vector  $\begin{pmatrix} x \\ y \end{pmatrix}$  in  $\mathbf{Z}^2$  by discussing whether  $|x| < |y|$  etc. You might want to draw a picture.

Now prove that the subgroup  $\langle u, v \rangle$  generated by  $\{u, v\}$  is the free group on  $\{u, v\}$ .

*Hint: first, understand that this amounts simply to proving that every (non-empty) reduced word using  $u, v$  represents a non-trivial element of  $\mathrm{SL}_2(\mathbf{Z})$ .*

## Exercise 3

- (i) Prove that  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  generate  $\mathrm{SL}_2(\mathbf{Z})$  and satisfy a non-trivial relation.
- (ii) Deduce that  $\mathrm{SL}_2(\mathbf{Z})$  itself is not free.
- (iii) Prove that  $\mathrm{SL}_2(\mathbf{Z})$  is “virtually free” in the sense that the quotient  $\mathrm{SL}_2(\mathbf{Z})/\langle u, v \rangle$  is finite.

## Exercise 4\*

Prove the following:

If a sequence  $\{v_n\}_{n \in \mathbf{N}}$  of  $\ell^1(\mathbf{Z}) \subseteq \ell^\infty(\mathbf{Z})^*$  converges weak-\* to some  $v \in \ell^\infty(\mathbf{Z})^*$ , then  $v \in \ell^1(\mathbf{Z})$ .

(Notice that this statement implies in particular the result of Exercise 2 from Problem Set 2.)

*Why does this not contradict Exercise 2(iii) from Problem Set 1?!?*