

STRUCTURAL PROPERTIES OF DENDRITE GROUPS

BRUNO DUCHESNE AND NICOLAS MONOD

ABSTRACT. Let G be the homeomorphism group of a dendrite. We study the normal subgroups of G . For instance, there are uncountably many non-isomorphic such groups G that are simple groups. Moreover, these groups can be chosen so that any isometric G -action on any metric space has a bounded orbit. In particular they have the fixed point property (FH).

1. INTRODUCTION

This article investigates the class of groups $G = \text{Homeo}(X)$ that appear as homeomorphism group of a dendrite X , recalling that a *dendrite* is a locally connected continuum without simple closed curves.

Not much can be said unless we ask that the relationship between the group G and the dendrite X has some substance: for instance, there exists complicated dendrites with trivial homeomorphism group, and conversely the rather trivial “star” dendrite contains any countable group in its homeomorphism group.

We shall therefore focus on *dendro-minimal* dendrites, namely dendrites X that do not admit a proper sub-dendrite $Y \subseteq X$ invariant under $G = \text{Homeo}(X)$.

Our main goal is to relate the properties of G , as an abstract group, with those of the topological space X . Inbetween these two worlds, we can also consider G as a topological transformation group and hence as a Polish group.

We start with some evidence that this whole enterprise has content:

Theorem 1.1. *There is a family of 2^{\aleph_0} dendro-minimal dendrites such that the corresponding groups are pairwise non-isomorphic simple groups.*

We prove Theorem 1.1 with the explicit family of *generalised Ważewski dendrites* D_S , where S is an arbitrary non-empty subset of $\{3, 4, \dots, \infty\}$. These classical dendrites are characterised by the fact that the order of every branch point lies in S and that such branch points are arcwise dense in D_S for every order in S .

For the non-isomorphy statement of Theorem 1.1, we shall prove that $\text{Homeo}(D_S)$, as a group, determines the set S . This can also be deduced from Rubin’s deep Reconstruction Theorem [Rub89] but we provide a direct proof (see Section 7 for a discussion).

By contrast, the simplicity statement of Theorem 1.1 results from a general analysis of the normal subgroups of $G = \text{Homeo}(X)$ for all dendro-minimal dendrites X . A first distinction to make is whether X contains a *free arc*, namely an embedded copy $I \cong [0, 1]$ of an interval whose interior is open in X . This produces a copy in G of the orientation-preserving group

Date: October 2016.

B.D. is supported in part by French projects ANR-14-CE25-0004 GAMME and ANR-16-CE40-0022-01 AGIRA.

$\text{Homeo}_+([0, 1])$ and it turns out that G can be decomposed as a permutational wreath product over the set $\text{Free}(X)$ of all maximal free arcs, see Section 3 for exact definitions for the following:

Proposition 1.2. *Let X be a dendro-minimal dendrite. Suppose that X admits some free arc (but is not reduced to an arc). Then*

$$\text{Homeo}(X) \cong \prod_{I \in \text{Free}(X)} \text{Homeo}_+(I) \rtimes \text{Aut}(\text{Free}(X)).$$

For the analysis of the normal subgroups of $G = \text{Homeo}(X)$ with X general, consider the normal subgroup $G_{\text{Ends}} \triangleleft G$ generated by all stabilisers of end points of X , and likewise $G_{\text{Br}} \triangleleft G$ the group generated by all stabilisers of branch points.

Theorem 1.3. *Let X be any dendro-minimal dendrite not reduced to an arc. Then $G_{\text{Ends}} = G$. Moreover, any subgroup normalised by G_{Br} either contains G_{Br} or fixes $\text{Ends}(X)$ pointwise.*

This will imply the following simplicity statements.

Corollary 1.4. *Let X be any dendro-minimal dendrite.*

If X has no free arc, then G_{Br} is a simple group.

If X admits some free arc (but is not reduced to it), then the image of G_{Br} in $\text{Aut}(\text{Free}(X))$ is a simple group.

The approach of the present article is in a sense opposite to our approach in [DM16], where we studied obstructions for group actions on dendrites. There, we took the external view-point of representing some other group into G . Several of these results pointed to analogies with negative curvature, especially with results for isometric actions on trees. However, to stress how different the topological setting of dendrites can be, we proposed the following problem:

Find a Kazhdan group with a non-elementary action on a dendrite.

Such a group would have a fixed point for any isometric action on a tree, or on a complete \mathbf{R} -tree.

Currently we do not have such an example if *Kazhdan group* is understood in the context of abstract groups, which must then be finitely generated. We do however have many examples with *property (FH)*, i.e. the fixed-point property for isometric actions on Hilbert spaces. The latter property is precisely the consequence of Kazhdan's property which forbids actions on trees and on \mathbf{R} -trees. It is moreover equivalent to Kazhdan's property for countable groups.

In fact the next result is much stronger than property (FH) and shows that the dendrites D_S are strongly allergic to the very idea of any metric structure.

Theorem 1.5. *Let S be a non-empty subset of $\{3, 4, \dots, \infty\}$. Then any isometric action of $\text{Homeo}(D_S)$ on any metric space has a bounded orbit.*

In particular, the group $\text{Homeo}(D_S)$ has property (FH).

Another way to come close to an answer to the problem of Kazhdan groups is to consider $\text{Homeo}(X)$ with its natural topology as a homeomorphism group. We can then use a result of Evans–Tsankov [ET16] and deduce that certain dendrite groups even have the *strong* version of Kazhdan's property, namely admit a finite Kazhdan set.

Theorem 1.6. *Let S be a finite non-empty subset of $\{3, 4, \dots, \infty\}$.*

Then the Polish group $\text{Homeo}(D_S)$ has the strong Kazhdan property (T).

Location of the proofs. The simplicity proof for Theorem 1.1 is completed in Corollary 6.4 and the non-isomorphy is Corollary 7.4. Proposition 1.2 is contained in Theorem 3.4. Theorem 1.3 follows from Theorems 4.1 and 4.4. Corollary 1.4 is then deduced as Corollaries 4.5 and 4.6. Theorem 1.5 occurs as Corollary 6.12 below, and Theorem 1.6 as Corollary 6.9.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Normal subgroups from free arcs	7
4. Normal subgroups from stabilisers	9
5. Semi-linear orders	12
6. Generalized Ważewski dendrites	16
7. Non-isomorphic homeomorphism groups	20
References	23

2. PRELIMINARIES

A general reference for dendrites is §10 in [Nad92]. We shall also use the basic properties of *elementary* and *dendro-minimal* actions on dendrites as discussed in [DM16]. We now recall some of the background but refer to the above references for the points not justified below.

2.A. Dendrites. In the case of a dendrite X , the Menger–Urysohn *order* of a point $x \in X$ is simply the cardinality of the set of components of $X \setminus \{x\}$. The points of order one, two or ≥ 3 form respectively the sets $\text{Ends}(X)$ of *end points*, $\text{Reg}(X)$ of *regular points* and $\text{Br}(X)$ of *branch points*. For $3 \leq n \leq \infty$, we write $\text{Br}_n(X)$ for the set of points of order n . The set $\text{Br}(X)$ is always countable, $\text{Reg}(X)$ is arcwise dense and uncountable, and $\text{Ends}(X)$ is non-empty (assuming X is not reduced to a point).

If $Y \subseteq X$ is a non-empty closed connected subset (thus a sub-dendrite), then there exists a canonical continuous retraction $X \rightarrow Y$ called the *first-point map*.

Any pair of points x, y is connected by a unique topological arc in X , denoted $[x, y]$. A *free arc* is an arc, not reduced to a point, whose interior is open in X . Since X is a dendrite, it is equivalent to ask that the interior of the arc contains no branch point. Any free arc is contained in a maximal free arc; we denote by $\text{Free}(X)$ the set of all maximal free arcs of X , which is a countable set. Two maximal free arcs meet at most at a common extremity, which is then a branch point. The following are equivalent, provided X is not reduced to a point: X has no free arc; $\text{Br}(X)$ is dense in X ; $\text{Ends}(X)$ is dense in X .

Any continuous self-map of a dendrite has a fixed point. A non-trivial arc $I = [x, y]$ in X is *austro-boreal* for a given homeomorphism $g \in \text{Homeo}(X)$ if the intersection of the fixed-point set of g in X with I is $\{x, y\}$. In that case, g preserves I and its action on $I \setminus \{x, y\}$ is conjugated to a translation action on \mathbf{R} .

The following lemma can be deduced e.g. from [KW30, §12].

Lemma 2.1. *Let d be a metric inducing the topology of the dendrite X . For every $\epsilon > 0$, there is a finite set F such that the components of $X \setminus F$ have d -diameter less than ϵ . \square*

It could alternatively be deduced from the following special case of [Why42, V.2.6], which we shall also need below.

Lemma 2.2. *Any sequence of disjoint connected subsets of a dendrite is a null-sequence, i.e. the diameter of its members tends to zero for any compatible metric.* \square

Remark 2.3. The set F of Lemma 2.1 can moreover be taken to belong to any given arcwise dense subset of X . Indeed, if F_0 is as in the lemma but for some $\epsilon_0 < \epsilon/2$, then any set F containing a point in $[x, y]$ for all distinct $x, y \in F_0$ will satisfy the condition for ϵ . On the other hand, one cannot restrict F to lie in a subset that is merely dense, as evidenced by dendrites with a dense set of ends.

2.B. Topologies on $\text{Homeo}(X)$. A natural topology on the homeomorphism group $\text{Homeo}(X)$ of a dendrite X is the topology of uniform convergence. With this topology, $\text{Homeo}(X)$ is a Polish group (see e.g. [Kec95, §9.B.9]). When X has no free arcs, a homeomorphism is completely determined by its action on $\text{Br}(X)$ since $\text{Br}(X)$ is dense in X . In particular, there is a faithful representation $\text{Homeo}(X) \rightarrow \text{Sym}(\text{Br}(X))$, where Sym is used to denote the group of all permutations of a given set. Since the set of branch points $\text{Br}(X)$ is countable, $\text{Sym}(\text{Br}(X))$ is a Polish group for the pointwise convergence. This gives a priori two topologies on $\text{Homeo}(X)$; in fact they coincide.

Proposition 2.4. *If X is a dendrite without free arcs, the homomorphism $\text{Homeo}(X) \rightarrow \text{Sym}(\text{Br}(X))$ is a homeomorphism onto its image.*

In particular, $\text{Homeo}(X)$ embeds as a closed subgroup of $\text{Sym}(\text{Br}(X))$.

Since $\text{Homeo}(X)$ is a closed subgroup of $\text{Sym}(\text{Br}(X))$, it implies in particular that it is a totally disconnected group; more precisely, its open subgroups form a neighbourhood system for the identity.

Proof. We first show that the topology induced from $\text{Sym}(\text{Br}(X))$ is finer than the uniform topology. The pointwise stabiliser in $\text{Homeo}(X)$ of a finite set $F \subseteq \text{Br}(X)$ as in Lemma 2.1 permutes the components of $X \setminus F$. Let $F' \supseteq F$ be a finite set containing also a point of each of these components. Then the pointwise stabiliser of F' preserves each connected component of $X \setminus F$ and therefore is ϵ -close to the identity in the uniform metric.

Conversely, let (g_n) be a sequence in $\text{Homeo}(X)$ converging to the identity in the uniform topology. For any $b \in \text{Br}(X)$, we choose x_1, x_2, x_3 in distinct components of $X \setminus \{b\}$. Since the connected components of $X \setminus \{b\}$ are open, we know that $g_n(x_i)$ and x_i belongs to the same component for n large enough. Since b is the only point in $[y_1, y_2] \cap [y_2, y_3] \cap [y_3, y_1]$ for any choice of elements y_i in the corresponding components, $g_n(b) = b$ follows.

Finally, the image of $\text{Homeo}(X)$ in $\text{Sym}(\text{Br}(X))$ is closed because it is Polish, see Exercise 9.6 in [Kec95]. \square

2.C. Separating subsets. The following elementary disjunction is Lemma 4.3 in [DM16].

Lemma 2.5. *Let G be a group with a dendro-minimal action on a dendrite X . For any proper sub-dendrite $Y \subseteq X$ there is $g \in G$ such that $gY \cap Y = \emptyset$.* \square

The proof given in [DM16] was as follows: if $gY \cap Y$ were never empty, then by a basic Helly theorem the intersection of gY over all $g \in G$ would be non-empty, contradicting dendro-minimality.

We shall use the following strengthening of Lemma 2.5.

Lemma 2.6. *Let G be a group with a dendro-minimal action on a dendrite X .*

For any proper sub-dendrites $Y, Y' \subseteq X$ there is $g \in G$ such that $gY \cap Y' = \emptyset$.

Proof. We can assume that Y meets Y' since otherwise $g = e$ will do. Therefore, $Y \cup Y'$ is a sub-dendrite of X . If this sub-dendrite is proper, then Lemma 2.5 applied to $Y \cup Y'$ provides g with $g(Y \cup Y') \cap (Y \cup Y') = \emptyset$ and a fortiori $gY \cap Y' = \emptyset$. Therefore we can assume $Y \cup Y' = X$. Now Lemma 2.5 yields g_1 with $g_1Y \cap Y = \emptyset$ and hence $g_1Y \subseteq Y'$. One last application of that lemma gives us g_2 with $g_2Y' \cap Y' = \emptyset$. We conclude that $g = g_2g_1$ satisfies $gY \subseteq g_2Y' \subseteq X \setminus Y'$ as was to be shown. \square

A minor variation of Lemma 2.6 is as follows.

Corollary 2.7. *Let G be a group with a dendro-minimal action on a dendrite X .*

For any connected set $Z \subseteq X$ that is not dense, any point $p \in X$ and any component C of $X \setminus \{p\}$, there is $g \in G$ such that $gZ \subseteq C$.

Proof. The complement $Y' = X \setminus C$ is a proper sub-dendrite and we can therefore apply Lemma 2.6 with $Y = \overline{Z}$. \square

We record a more technical variant of Corollary 2.7.

Lemma 2.8. *Let G be a group acting dendro-minimally on a dendrite X . Let $I \subseteq X$ be an arc and b be a branch point in the interior of I . Let $x \in X$ and let Y be a component of $X \setminus \{x\}$.*

Then there is $g \in G$ with $gI \subseteq Y$ such that the image of x under the first-point map to gI is gb .

Proof. Upon replacing g by g^{-1} , the task is to find $g \in G$ such $I \subseteq gY$ and b is the retraction of gx on I . Let p be a regular point on an arc branching off from I at b , let C be the component of $X \setminus \{p\}$ not containing b and denote by Z the dendrite $\{x\} \cup (X \setminus Y)$. Then it suffices to have $gZ \subseteq C$, which is possible by Corollary 2.7. \square

2.D. Patchwork. The following patchwork for homeomorphisms fails already if we replace dendrites by a disc.

Lemma 2.9. *Let \mathcal{U} be a family of disjoint open connected subsets of a dendrite X and let $(f_U)_{U \in \mathcal{U}}$ be a family of homeomorphisms $f_U \in \text{Homeo}(U)$ for $U \in \mathcal{U}$. Suppose that each f_U can be extended continuously to the closure \overline{U} by the identity on the boundary $\overline{U} \setminus U$.*

Then the map $f: X \rightarrow X$ given by f_U on each $U \in \mathcal{U}$ and the identity everywhere else is a homeomorphism.

Proof. By compactness, it suffices to prove that there cannot be a sequence $x_n \rightarrow x$ in X such that $f(x_n)$ converges to some $y \neq f(x)$. If there is such a sequence, then each $U \in \mathcal{U}$ contains x_n for at most finitely many indices n . Moreover, upon extracting we can assume that each x_n belongs to some $U_n \in \mathcal{U}$. By the previous observation, we can assume that all U_n are distinct and hence they form a null-sequence by Lemma 2.2. Since $f(x_n) = f_{U_n}(x_n)$ belongs also to U_n , we conclude that $f(x_n)$ converges to $f(x)$, contrary to our assumption. \square

2.E. Uniform structure. Like any compact space, a dendrite admits a unique compatible uniform structure.

Lemma 2.10. *For any dendrite X , the sets*

$$U_F = \{(x, y) \in X^2 : |[x, y] \cap F| \leq 1\}$$

form a basis of the uniform structure as F ranges over all finite subsets of X . Moreover, one can restrict F to belong to any arcwise dense subset of X .

Proof. These sets belong to the uniformity associated to the compact topology, i.e. they are neighbourhoods of the diagonal in $X \times X$: this follows from the fact that X is locally arcwise connected (like any Peano continuum [Nad92, 8.25]). In order to prove that they form a basis, choose a compatible metric d on X and $\epsilon > 0$. Let F be a finite set as provided by Lemma 2.1; we can assume that F is in a given arcwise dense subset by Remark 2.3. If now $(x, y) \in U_F$, then $[x, y] \setminus F$ has at most two components and thus $d(x, y) \leq 2\epsilon$. \square

2.F. Permutational wreath products. We begin with a result about Cartesian products.

Lemma 2.11. *Suppose that the direct product $P = \prod_{N \in \mathcal{N}} N$ of a family \mathcal{N} of groups acts on a dendrite. If each $N \in \mathcal{N}$ has fixed points, then so does P .*

Proof. By a compactness argument, we can choose a minimal P -invariant sub-dendrite $Y \subseteq X$. Each N has fixed points in Y since the first-point map $X \rightarrow Y$ is P -equivariant by its naturality. Since P normalises N it follows that $[Y \cap \text{Fix}(N)]$ is a P -invariant sub-dendrite and hence $[Y \cap \text{Fix}(N)] = Y$. It follows that $\text{Ends}(Y)$ is contained in $\text{Fix}(N)$ by [DM16, Lemma 2.3]. Since this holds for each $N \in \mathcal{N}$, we conclude that P fixes $\text{Ends}(Y)$. \square

Proposition 2.12. *Let N be a group without index two subgroup and let \mathcal{C} be a set of cardinality $3 \leq |\mathcal{C}| \leq \aleph_0$. Then every action of the full permutational wreath product*

$$H = N^{\mathcal{C}} \rtimes \text{Sym}(\mathcal{C})$$

on a dendrite has a fixed point.

Proof. Let X be a dendrite with an H -action. Since $N^{\mathcal{C}}$ is a product, one of its factors must act elementarily on X by Corollary 4.6 in [DM16] (that reference considers products of two factors, but the general case follows at once). Since N has no index two subgroup, elementarity means that this factor fixes a point in X . Since all factors of $N^{\mathcal{C}}$ are conjugated in H , they all fix some point. Therefore, $N^{\mathcal{C}}$ has a fixed point by Lemma 2.11.

We consider first the case where \mathcal{C} has finite size $n \geq 3$. Then H has a finite orbit and hence acts elementarily. If H does not fix a point, it preserves some arc $I \subseteq X$ and admits an index two subgroup H' preserving the orientation of I . We claim that H' is $N^{\mathcal{C}} \rtimes \text{Alt}(n)$. Since N has no index two subgroup, it suffices to check that $\text{Alt}(n)$ is the only index two subgroup of $\text{Sym}(n)$, or equivalently that it has no index two subgroup itself. Indeed, for $n \neq 4$ it is simple (of order $\neq 2$) and for $n = 4$ its only proper normal subgroup has index three. This proves the claim.

Since the group of orientation-preserving homeomorphisms of an interval contains no non-trivial finite subgroup, the image of $\text{Alt}(n)$ in $\text{Homeo}_+(I)$ is trivial. There is an element $\sigma \in \text{Sym}(n)$ reversing the orientation of I ; note that σ has a necessarily unique fixed point p in the interior of I . It suffices to prove that every factor $N_1 \cong N$ of $N^{\mathcal{C}}$ fixes p . If not, then for some $g \in N_1$ we have $gp > p$ with respect to a choice of an orientation of I . Then $\sigma g \sigma^{-1} p = \sigma gp < p$. On the other hand, $\text{Alt}(n)$ acts transitively on \mathcal{C} since $n \geq 3$. Therefore there is $\tau \in \text{Alt}(n)$ with $\tau g \tau^{-1} = \sigma g \sigma^{-1}$. It follows $\tau g \tau^{-1} p < p$, which contradicts $gp > p$ since τ acts trivially on I .

We now consider the case where \mathcal{C} is infinite countable. Denote by $\text{Sym}_f(\mathcal{C})$ the normal subgroup of finitely supported permutations of \mathcal{C} and by $\text{Alt}_f(\mathcal{C})$ its alternating subgroup. Since $\text{Sym}_f(\mathcal{C})$ is an amenable group and since $N^{\mathcal{C}}$ fixes a point, Lemma 6.1 in [DM16] implies that $N^{\mathcal{C}} \rtimes \text{Sym}_f(\mathcal{C})$ acts elementarily. Let thus $I \subseteq X$ be an invariant arc; note that $N^{\mathcal{C}}$ fixes a point of I , namely the image of any fixed point under the first-point map

$X \rightarrow I$. As above, $\text{Alt}_f(\mathcal{C})$ acts trivially on I , being an increasing union of $\text{Alt}(n)$. Therefore, $N^{\mathcal{C}} \rtimes \text{Alt}_f(\mathcal{C})$ fixes a point. The non-empty set $K \subseteq X$ of points fixed by $N^{\mathcal{C}} \rtimes \text{Alt}_f(\mathcal{C})$ is closed, and therefore it is a compact metrisable space. It is H -invariant and the H -action on K descends to an action of $\text{Sym}(\mathcal{C}) / \text{Alt}_f(\mathcal{C})$. If we endow $\text{Sym}(\mathcal{C})$ with its Polish topology, the subgroup $\text{Alt}_f(\mathcal{C})$ is dense. However, Kechris–Rosendal have proved that any action of $\text{Sym}(\mathcal{C})$ on a compact metrisable space is continuous for this topology, see Theorem 1.10 in [KR07]. Therefore, the $\text{Sym}(\mathcal{C})$ action on K is trivial and H fixes every point of K . \square

We record the following for later use.

Lemma 2.13. *Let N be any group and let \mathcal{C} be an infinite countable set. Then the group*

$$H = N^{\mathcal{C}} \rtimes \text{Sym}(\mathcal{C})$$

has no index two subgroup.

Proof. Schreier–Ulam proved in [SU33, §1.7] that the only proper normal subgroups of $\text{Sym}(\mathcal{C})$ are $\text{Sym}_f(\mathcal{C})$ and $\text{Alt}_f(\mathcal{C})$. Therefore, an index two subgroup of H would have to be of the form $M \rtimes \text{Sym}(\mathcal{C})$, where $M \triangleleft N^{\mathcal{C}}$ is of index two and $\text{Sym}(\mathcal{C})$ -invariant. In particular, for every $y \in N^{\mathcal{C}}$ and every $\sigma \in \text{Sym}(\mathcal{C})$, we have $y^\sigma y^{-1} \in M$. Let now z be an arbitrary element of $N^{\mathcal{C}}$. It suffices to express z as $y^\sigma y^{-1}$. To this end, we take σ to be the shift under identifying \mathcal{C} with \mathbf{Z} . We can now define $y = (y_n)_{n \in \mathbf{Z}}$ inductively by taking y_0 arbitrary and setting $y_n = z_{n-1} y_{n-1}$ for $n > 0$ and $y_n = z_n^{-1} y_{n+1}$ for $n < 0$. \square

2.G. The generalised Ważewski dendrites D_S . Let S be a non-empty subset of $\{3, 4, \dots, \infty\}$; here ∞ , standing for “not finite”, will in fact be the cardinal \aleph_0 . There exists a dendrite, not reduced to a point, whose branch points all have orders in S and with the following property: for every $n \in S$, every arc contains an element of order n . It follows that $\text{Br}_n(D_S)$ is arc-wise dense. The existence of such a dendrite can be proved for instance by an inverse limit construction following the procedure given in [Nad92, 10.37] for the special case $S = \{\infty\}$. In fact, such a dendrite is unique up to homeomorphism by Theorem 6.2 in [CD94]. It is called the *generalised Ważewski dendrite D_S* . The uniqueness implies:

Lemma 2.14. *Every non-trivial sub-dendrite of D_S is homeomorphic to D_S .* \square

We point out that $\text{Ends}(D_S)$ is uncountable. Indeed, it has no isolated points and it is Polish since it is a dense G_δ in D_S . The latter is the case because its complement can be written as the union of all arcs $[x, y]$ where x, y range over the countable set $\text{Br}(D_S)$.

3. NORMAL SUBGROUPS FROM FREE ARCS

There are several natural normal subgroups of $\text{Homeo}(X)$ related to free arcs: we define

$$\text{Fix}(\text{Free}(X)) < \text{Stab}_+(\text{Free}(X)) < \text{Stab}(\text{Free}(X))$$

to consist of the homeomorphisms for which every $I \in \text{Free}(X)$ is fixed pointwise, respectively preserved together with its orientation, respectively just preserved.

Furthermore, we denote by $\text{Fix}(\text{Ends}(X))$ the normal subgroup of $\text{Homeo}(X)$ consisting of the elements that fix every end of X .

Proposition 3.1. *Let X be a dendrite admitting some free arc.*

Then the group $\text{Fix}(\text{Ends}(X))$ is contained in $\text{Stab}_+(\text{Free}(X))$ and we have a decomposition

$$\text{Stab}_+(\text{Free}(X)) = \text{Fix}(\text{Free}(X)) \times \text{Fix}(\text{Ends}(X)).$$

Moreover, the restriction maps to $\text{Homeo}(I)$ induce an isomorphism

$$\text{Fix}(\text{Ends}(X)) \cong \prod_{I \in \text{Free}(X)} \text{Homeo}_+(I).$$

Proof. We claim that $\text{Fix}(\text{Ends}(X))$ fixes every branch point p of X . Indeed, there exists three arcs meeting only at p ; by a maximality argument, the other extremity of each of those can be taken to be an end of X . Any homeomorphism fixing these three ends fixes p , whence the claim. Since any extremity of a maximal free arc is either an end, or a branch point, or a limit of branch points, the claim shows already that $\text{Fix}(\text{Ends}(X))$ is contained in $\text{Stab}_+(\text{Free}(X))$.

Next, we claim that the two normal subgroups $\text{Fix}(\text{Ends}(X))$ and $\text{Fix}(\text{Free}(X))$ intersect trivially; hence they commute and generate their direct product. For this we need to show that any homeomorphism in their intersection fixes every regular point of X . This is the case by the previous claim since any regular point either is a limit of branch points or is contained in a free arc.

Finally, we consider the restriction morphism from $\text{Stab}_+(\text{Free}(X))$ to $\prod_{I \in \text{Free}(X)} \text{Homeo}_+(I)$, whose kernel is exactly $\text{Fix}(\text{Free}(X))$; in particular it is injective on $\text{Fix}(\text{Ends}(X))$. On the other hand this map is surjective when restricted to $\text{Fix}(\text{Ends}(X))$ because of Lemma 2.9 applied to the interiors of all $I \in \text{Free}(X)$. Both isomorphisms of Proposition 3.1 follow. \square

Proposition 3.2. *Let X be a dendrite admitting some free arc.*

If X is dendro-minimal, then the union of all maximal free arcs is dense in X .

Proof. It suffices to prove that any branch point $p \in X$ is a limit of points belonging to free arcs. Choose $q \neq p$; we can assume that $[p, q]$ contains a sequence (p_n) of distinct branch points p_n converging to p because otherwise p would bound a free arc in $[p, q]$. Let c_n be a component of $X \setminus \{p_n\}$ not containing p nor q . Then all c_n are pairwise disjoint and therefore form a null-sequence by Lemma 2.2. It remains only to observe that each c_n contains a free arc; this follows from Corollary 2.7, choosing Z to be some free arc. \square

Corollary 3.3. *Let X be a dendrite admitting some free arc. If X is dendro-minimal, then*

$$\text{Stab}_+(\text{Free}(X)) = \text{Fix}(\text{Ends}(X)) \cong \prod_{I \in \text{Free}(X)} \text{Homeo}_+(I).$$

Furthermore,

$$\text{Stab}(\text{Free}(X)) = \text{Stab}_+(\text{Free}(X))$$

unless X is reduced to an arc.

Proof. By Proposition 3.2, the group $\text{Fix}(\text{Free}(X))$ is trivial. Therefore, the first statement follows from Proposition 3.1.

It remains to justify that every element g of $\text{Stab}(\text{Free}(X))$ preserves the orientation of any $I \in \text{Free}(X)$ unless $X = I$. Thus we can assume that at least one extremity p of I is not an end of X . Therefore, there is a component c of $X \setminus \{p\}$ which does not meet I . By Corollary 2.7, c contains a maximal free arc J , namely an image of I . Since g preserves J and I , it cannot permute the extremities of I . \square

Define $\text{Aut}(\text{Free}(X))$ to be the image of $\text{Homeo}(X)$ in the permutation group of the countable set $\text{Free}(X)$; in other words,

$$\text{Aut}(\text{Free}(X)) = \text{Homeo}(X) / \text{Stab}(\text{Free}(X)).$$

Theorem 3.4. *Let X be a dendrite admitting some free arc, but not reduced to an arc.*

If X is dendro-minimal, then $\text{Homeo}(X)$ admits a semi-direct product decomposition

$$\text{Homeo}(X) \cong \text{Fix}(\text{Ends}(X)) \rtimes \text{Aut}(\text{Free}(X))$$

and hence it is the permutational wreath product

$$\text{Homeo}(X) \cong \prod_{I \in \text{Free}(X)} \text{Homeo}_+(I) \rtimes \text{Aut}(\text{Free}(X)).$$

Proof. We choose for each $I \in \text{Free}(X)$ a homeomorphism $\alpha_I: [0, 1] \rightarrow I$. We further choose an involution τ of $[0, 1]$ that exchanges the end-points.

If now g is an arbitrary element of $\text{Homeo}(X)$, we define for $I \in \text{Free}(X)$ the involution τ_I^g of $[0, 1]$ to be trivial if $g\alpha_I$ and α_{gI} give the same orientation to gI and to be τ otherwise. This can be written more obscurely as $\tau_I^g = \tau^{\alpha_{gI}^{-1}g\alpha_I(0)}$; in any case, we have the cocycle relation $\tau_I^{g_1g_2} = \tau_{g_2I}^{g_1}\tau_I^{g_2}$. We define furthermore the map f_I^g on I by $f_I^g = \alpha_I \circ \tau_I^g \circ \alpha_{gI}^{-1} \circ g|_I$. Then f_I^g is in $\text{Homeo}_+(I)$. Therefore, Proposition 3.1 ensures that there is $f^g \in \text{Fix}(\text{Ends}(X))$ with $f^g|_I = f_I^g$ for all $I \in \text{Free}(X)$. We can therefore define an element $\bar{g} \in \text{Homeo}(X)$ by $\bar{g} = g \circ (f^g)^{-1}$.

The cocycle relation for τ_I^g shows that the assignment $g \mapsto \bar{g}$ is a group homomorphism. More precisely, this holds by construction when we evaluate on any point in a free arc and thus everywhere since the union of all free arcs is dense by Proposition 3.2.

This homomorphism is trivial on $\text{Stab}(\text{Free}(X))$. Indeed, any g in that group preserves the orientation of every free arc since X is not an arc, see Corollary 3.3. Therefore the construction of f_I^g shows that $f_I^g = g|_I$, hence \bar{g} is trivial. In conclusion, the morphism $g \mapsto \bar{g}$ provides a section from $\text{Aut}(\text{Free}(X))$ to $\text{Homeo}(X)$ because f^g is in $\text{Fix}(\text{Ends}(X))$ which coincides with $\text{Stab}(\text{Free}(X))$ by Corollary 3.3.

Finally, the wreath product statement follows from the naturality of the isomorphism between $\text{Fix}(\text{Ends}(X))$ and $\prod_{I \in \text{Free}(X)} \text{Homeo}_+(I)$. \square

4. NORMAL SUBGROUPS FROM STABILISERS

Let X be an arbitrary dendrite. A priori there is a number of natural normal subgroups of $\text{Homeo}(X)$ associated to point stabilisers. Just to keep the arguments below more legible, we temporarily write $G = \text{Homeo}(X)$ and introduce the following notation: G_{Br} , G_{Ends} and G_{Reg} are the subgroups of G generated by all stabilisers of branch points, respectively of end points and regular points. Similarly, G_{Arc} denotes the subgroup generated by all pointwise stabilisers of non-trivial arcs, and G_{Comp} by all pointwise stabilisers of connected components of the complement of points.

All these are closed normal subgroups of G .

Theorem 4.1. *Let X be any dendro-minimal dendrite not reduced to an arc. Then*

$$G_{\text{Arc}} = G_{\text{Br}} = G_{\text{Comp}} < G_{\text{Reg}} < G_{\text{Ends}} = G.$$

In fact, the first two equalities above are basic and more general:

Lemma 4.2. *For any dendrite not reduced to a point, $G_{\text{Arc}} = G_{\text{Br}} = G_{\text{Comp}} < G_{\text{Ends}} \cap G_{\text{Reg}}$.*

Proof of Lemma 4.2. By definition, we have $G_{\text{Comp}} < G_{\text{Arc}} \cap G_{\text{Br}} \cap G_{\text{Ends}} \cap G_{\text{Reg}}$. We first claim that $G_{\text{Br}} < G_{\text{Comp}}$. Let thus $g \in G$ fix a branch point x . We define a homeomorphism h that

coincides with g on some component C of $X \setminus \{x\}$, that coincides with g^{-1} on gC and is the identity elsewhere. Since $X \setminus \{x\}$ has at least three connected components, $h \in G_{\text{Comp}}$. Now, $g = (gh^{-1})h$ and gh^{-1} fixes C pointwise. Therefore, $g \in G_{\text{Comp}}$. This proves the first claim and hence $G_{\text{Br}} = G_{\text{Comp}}$.

Our second claim is that any element g fixing a regular point r without exchanging the two components C_1, C_2 of $X \setminus \{r\}$ belongs to G_{Comp} . Indeed, g can then be written $g = g_1 g_2$ where g_i fixes pointwise C_i and acts as g does on the other component.

This second claim implies in particular $G_{\text{Arc}} < G_{\text{Comp}}$ and thus $G_{\text{Arc}} = G_{\text{Comp}}$. \square

Proof of Theorem 4.1. We claim that any element $g \in G$ which admits an austro-boreal arc belongs to G_{Ends} . Indeed, this is the case if an extremity of such an arc is an end of X , but also if it is a branch point since $G_{\text{Br}} < G_{\text{Ends}}$ by Lemma 4.2. If on the other hand the extremities are regular points, then g cannot exchange the corresponding components since it preserves the arc. Thus we conclude again $g \in G_{\text{Comp}} < G_{\text{Ends}}$, this time by the second claim of the proof of Lemma 4.2.

Our main goal is to prove $G_{\text{Reg}} < G_{\text{Ends}}$. It suffices to consider an element $g \in G$ which fixes a regular point r and exchanges the components C_1, C_2 of $X \setminus \{r\}$. By Theorem 10.5 in [DM16], G contains an element h admitting an austro-boreal arc I . By Lemma 2.8, we can assume, after conjugating h , that I lies in C_2 and that the first-point map sends r to some branch point b in the interior of I .

We shall now prove that $hg \in G_{\text{Ends}}$, which implies indeed $g \in G_{\text{Ends}}$ since $h \in G_{\text{Ends}}$ by the above claim. We know that hg fixes some point $p \in X$. Given what we have established so far, the only problematic case is if p is a regular point and if moreover hg permutes the components of $X \setminus \{p\}$. We shall prove by contradiction that this cannot happen.

Indeed, in that case p is the only fixed point of hg and, moreover, for every $x \in X$ the arc $[x, hgx]$ contains p . We first apply this observation to $x = hb \in I$. Then the arc $[x, hgx]$ starts from hb , crosses hr and ends in hC_1 . Our assumption on the respective position of I and r shows that this entire arc projects to hb under the first-point map $X \rightarrow I$. We now apply the observation to $x = (hg)^{-1}b \in C_1$. This time the arc $[x, hgx]$ starts in C_1 , crosses r and ends at b . Therefore, this arc projects to b under the first-point map $X \rightarrow I$. It follows that the two arcs do not intersect, a contradiction.

At this point we know that both G_{Br} and G_{Reg} are contained in G_{Ends} ; since any homeomorphism of X must fix some point, it follows that $G = G_{\text{Ends}}$. \square

Lemma 4.3. *Let X be a dendrite and let N, H be subgroups of $\text{Homeo}(X)$.*

Suppose that H acts dendro-minimally on X and normalises N . Then either N acts dendro-minimally on X or N fixes $\text{Ends}(X)$ pointwise.

In particular, if X has no free arc, then any non-trivial normal subgroup of a dendro-minimal subgroup of $\text{Homeo}(X)$ is dendro-minimal as well.

Proof. If N has no fixed point in X , then there is a *unique* minimal N -invariant sub-dendrite Y in X , see Remark 4.2 in [DM16]. Since H normalises N , it preserves Y and thus $Y = X$.

If on the other hand N has a fixed point $x \in X$, then the H -orbit of x consists of N -fixed points, and hence so does its closure. This closure contains $\text{Ends}(X)$ by Lemma 4.4 in [DM16].

The additional statement follows from the fact that $\text{Ends}(X)$ is dense in X if X has no free arc (see e.g. [DM16, 4.7]). \square

Theorem 4.4. *Let X be any dendro-minimal dendrite not reduced to an arc. Let $N < G = \text{Homeo}(X)$ be any subgroup normalised by G_{Br} .*

Then either N contains G_{Br} or N fixes $\text{Ends}(X)$ pointwise.

If X has no free arc, we deduce:

Corollary 4.5. *Let X be any dendro-minimal dendrite without free arc.*

Then G_{Br} is a simple group. □

If on the contrary X admits some free arc, then we appeal to Theorem 3.4 and deduce:

Corollary 4.6. *Let X be any dendro-minimal dendrite with a free arc but not reduced to this arc.*

Then the image $G_{\text{Br}} / \text{Fix}(\text{Ends}(X))$ of G_{Br} in $\text{Aut}(\text{Free}(X))$ is a simple group. □

The proof of Theorem 4.4 uses an idea of Tits from [Tit70], in the following setup. Let I be an arc of a dendrite X . For any point t of I , denote by $X_t(I)$ the preimage of t under the first-point map $X \rightarrow I$. Thus, if $X_t(I)$ is not reduced to $\{t\}$, then t is either a branch point or an extremity of I . For any space Y and any $t \in Y$, we denote by $\text{Homeo}_t(Y)$ the stabiliser of t in $\text{Homeo}(Y)$.

Lemma 4.7. *In the above notation, suppose we are given $h_t \in \text{Homeo}_t(X_t(I))$ for each $t \in I$. Then there is $h \in \text{Fix}(I)$ such that $h|_{X_t(I)} = h_t$ for all $t \in I$.*

Proof. Whenever $X_t(I)$ is not reduced to $\{t\}$, it is the closure of the set $X_t(I) \setminus \{t\}$, which is the union of the components of $X \setminus \{t\}$ that do not meet I . Therefore, the lemma is a particular case of Lemma 2.9. □

Proof of Theorem 4.4. Assume that N does not fix $\text{Ends}(X)$ pointwise. Two applications of Lemma 4.3 show that N acts dendro-minimally on X because N is normalised by G_{Br} which is normal in G . Since $G_{\text{Br}} = G_{\text{Comp}}$, it suffices to show that every $g \in G$ fixing pointwise a component Y of $X \setminus \{x\}$ for some $x \in X$ belongs to N .

We know from Theorem 10.5 in [DM16] that N contains an element n admitting and austro-boreal arc $I = [y, z]$. By Lemma 2.8, we can assume, upon conjugating n within N , that I lies in Y and that the image b of x under the first-point map to I is some branch point in the interior of I .

The proof will be complete if we express g as $g = [h, n] = hnh^{-1}n^{-1}$ for some element h of $G_{\text{Arc}} = G_{\text{Br}}$, since then $g = (hnh^{-1})n^{-1} \in N$.

This element is provided by Lemma 4.7 as follows. Recall that the n -action on $I \setminus \{y, z\}$ is free by definition of austro-boreal arcs. We set h_t to be the identity unless $t = n^k b$ for some $k \geq 0$. In that case, we set $h_{n^k b} = n^k g n^{-k}$, which defines indeed a homeomorphism of $X_{n^k b}(I)$ fixing $n^k b$. Now the identity $g = [h, n]$ holds by construction, finishing the proof. □

We close this section with an observation about individual homeomorphisms fixing an end point.

Lemma 4.8. *Let g be a homeomorphism of a dendrite X not reduced to a point.*

If g fixes an end point, then g fixes at least two points.

Proof. Let x be a g -fixed end point and let y be any other point. We denote by $z \in [x, y]$ the point at which $[x, y]$ branches away from $[x, gy]$. Now gz belongs to $[x, gy]$ and thus either $[x, gz] \subseteq [x, z]$ or $[x, z] \subseteq [x, gz]$. Upon replacing g by g^{-1} and y by gy , we may assume that we are in the second case. Then the closure of $\bigcup_{n \in \mathbb{N}} [x, g^n z]$ is a non-trivial arc whose end points are g -fixed. □

5. SEMI-LINEAR ORDERS

The goal of this section is to relate dendrites to the classical theory of semi-linear orders. This relation being fruitful in both directions, we investigate it slightly beyond what is strictly needed for the results of the introduction. We begin by recalling some standard notions about orders:

Let (T, \leq) be an ordered set. A subset is a *chain* if it is linearly (=totally) ordered. One writes $x \downarrow = \{y : y \leq x\}$ and $x \uparrow = \{y : y \geq x\}$. An order is *dense* if for all x, y with $x < y$ there is t with $x < t < y$. More generally, a subset $T_0 \subseteq T$ is *order-dense in T* if for all $x, y \in T$ with $x < y$ there is $t \in T_0$ with $x < t < y$. A subset $T_0 \subseteq T$ is *cointial in T* if for all $x \in T$ there is $y \in T_0$ with $y \leq x$.

Definition 5.1. An ordered set (T, \leq) is (*lower*) *semi-linear* if the following two axioms hold:

T is downwards directed: $\forall x, y \exists s : s \leq x, y$;
for all $x \in T$, the set $x \downarrow$ is a chain.

Observe that any cointial subset of a semi-linear order is still semi-linear.

The definitions are readily checked in the following example:

Example 5.2. Let X be a dendrite and $z \in X$ an end of X . Then the relation defined on X by

$$x \leq y \iff [z, x] \subseteq [z, y]$$

is a dense semi-linear order.

More generally, any subset $T \subseteq X$ that is arcwise dense (in the topological sense) is order-dense in (X, \leq) and is a semi-linear order. (Although T is not cointial in X if $z \notin T$, it is then cointial in $X \setminus \{z\}$, which is still a semi-linear order.)

The main purpose of this section is to construct and study a functor reverting Example 5.2: to any semi-linear order T , we shall associate a “completion” \widehat{T} containing it. Under natural assumptions, \widehat{T} will be a dendrite inducing the given order on T . See Remark 5.12 for the relation to known completions.

Definition 5.3. Let (T, \leq) be a semi-linear order. We define the ordered set (\widehat{T}, \leq) to be the set of all *full down-chains* of T endowed with the inclusion order (induced from the set of all subsets of T). Here a chain $C \subseteq T$ is a *down-chain* if $x \downarrow \subseteq C$ for all $x \in C$. It is called *full* if it contains its supremum or has no supremum in T .

Recall that for a subset S of an ordered set T , an element $s \in T$ is a *supremum* of S in T if it is an upper bound for S and if $s \leq s'$ for every upper bound $s' \in T$ for S . In particular, a supremum is unique if it exists; it is sometimes denoted by $\vee S$. We should however remember that the definition depends on T since S might not have a supremum anymore in a larger ordered set containing T . The infimum is defined dually.

Remark 5.4. The empty set has a supremum in a semi-linear order T if and only if T admits a (necessarily unique) minimal element t . In that case, $\{t\}$ is a unique minimal element of \widehat{T} . On the other hand, T admits no minimum if and only if $\emptyset \in \widehat{T}$. In that case, \emptyset is a unique minimal element of \widehat{T} .

Thus, in any case \widehat{T} admits a unique minimum; this will turn out to be our end z .

The definitions imply readily the following.

Lemma 5.5. *Let (T, \leq) be a semi-linear order. Then the map*

$$T \longrightarrow \widehat{T}, \quad x \longmapsto x \downarrow$$

is well-defined, injective, and realizes T as a subset of \widehat{T} with the same induced order. \square

Lemma 5.6. *Let (T, \leq) be a semi-linear order. Then (\widehat{T}, \leq) is a semi-linear order too.*

Proof. The ordered set \widehat{T} is downwards directed since it has a minimum by Remark 5.4. For the other axiom, we pick $C \in \widehat{T}$ and need to check that $C \downarrow$ is a chain in \widehat{T} . This holds because the collection of down-chains in any chain is linearly ordered by inclusion. \square

Recall that an ordered set is \vee -complete if every non-empty subset admits a supremum; \wedge -completeness is defined dually.

Proposition 5.7. *Let (T, \leq) be a semi-linear order. Then (\widehat{T}, \leq) is \wedge -complete.*

Moreover, any non-empty chain $\mathcal{C} \subseteq \widehat{T}$ admits a supremum in \widehat{T} . In particular, all maximal chains $\mathcal{C} \subseteq \widehat{T}$ are Dedekind-complete.

Proof. Let $\mathcal{C} \subseteq \widehat{T}$ be a non-empty subset. Then $E = \bigcap_{C \in \mathcal{C}} C$ is a down-chain in T . If E has a supremum $\vee E$ in T , the fact that each $C \in \mathcal{C}$ is a full down-chain implies $\vee E \in C$ and thus $\vee E \in E$. Therefore $E \in \widehat{T}$ and E is an infimum for \mathcal{C} in \widehat{T} .

Suppose that \mathcal{C} is moreover a chain. Consider $F = \bigcup_{C \in \mathcal{C}} C$; it is a down-chain in T . If F has no supremum in T , then $F \in \widehat{T}$ and F is a supremum for \mathcal{C} in \widehat{T} . If on the other hand F has a supremum $\vee F$ in T , let $F^* = F \cup \{\vee F\}$. Then $F^* \in \widehat{T}$ and F^* is an upper bound for \mathcal{C} . We claim that F^* is a supremum for \mathcal{C} in \widehat{T} . Indeed, let $F' \in \widehat{T}$ be an upper bound for \mathcal{C} . We need to show $F^* \subseteq F'$ and know already $F \subseteq F'$. Since F is not full, $F \neq F'$. Let thus $t \in F' \setminus F$. Then t is an upper bound for F since F' is a chain and F' a down-chain. Therefore $\vee F \leq t$ and since F' is a down-chain we deduce $\vee F \in F'$ as desired.

Finally, for Dedekind-completeness, it suffices to observe that the supremum and infimum constructed in T must automatically belong to any maximal chain containing \mathcal{C} . \square

In the converse direction, we have:

Lemma 5.8. *Let (T, \leq) be a semi-linear order. Suppose that every non-empty chain in T has a supremum and an infimum in T . Then the morphism $T \rightarrow \widehat{T}$ is an isomorphism.*

Proof. We need to prove that any $C \in \widehat{T}$ is of the form $C = x \downarrow$ for some $x \in T$. We begin by noting that $C \neq \emptyset$, or equivalently by Remark 5.4, that T has a minimum. Indeed, the infimum of any maximal chain in T is a minimum for T since T is downwards directed. Being non-empty, C has a supremum x in T . Being a full down-chain, C coincides with $x \downarrow$. \square

Remark 5.9. The above proof shows that Lemma 5.8 holds under the formally weaker assumption that every non-empty chain in T has a supremum in T and that T has a minimum.

Taking into account Proposition 5.7, we deduce:

Corollary 5.10. *Let (T, \leq) be a semi-linear order. Suppose that every non-empty chain in T has a supremum in T and that T has a minimum. Then (T, \leq) is \wedge -complete. \square*

Combining again Proposition 5.7 with Lemma 5.8, we also deduce that the operation $T \mapsto \widehat{T}$ is idempotent.

Corollary 5.11. *For any semi-linear order (T, \leq) , the morphism $\widehat{T} \rightarrow \widehat{\widehat{T}}$ is an isomorphism. \square*

Remark 5.12. The above results show that \widehat{T} is a completion of T in some sense. It is larger than the Dedekind-type completion of [Dro85, 5.3], [Mar89, 1.3] but smaller than the MacNeille completion [Mac37].

Example 5.13. Consider the subset $T = [-1, 0) \cup \{i, -i\}$ of the complex numbers endowed with the order given by the usual order on the real part (but i incomparable to $-i$). This is a dense semi-linear order. Moreover, \widehat{T} can be identified with the closure $T \cup \{0\}$ with the order extended by $0 \leq i, 0 \leq -i$. This semi-linear order is however not dense anymore.

The construction of \widehat{T} is functorial in the following sense.

Lemma 5.14. *Let (T, \leq) be a semi-linear order and $T_0 \subseteq T$ a subset. Suppose that (T_0, \leq) is semi-linear (e.g. it is coinital in T). Then there is a canonical inclusion of ordered sets $\widehat{T}_0 \rightarrow \widehat{T}$ making the diagram*

$$\begin{array}{ccc} T & \longrightarrow & \widehat{T} \\ \uparrow & & \uparrow \\ T_0 & \longrightarrow & \widehat{T}_0 \end{array}$$

commutative.

Proof. In view of Lemma 5.8 and Corollary 5.11, we can simplify the notation by assuming that T admits a supremum and infimum for each of its non-empty chains; we then construct a suitable map $\widehat{T}_0 \rightarrow T$. Namely, we associate to $C \in \widehat{T}_0$ its supremum in T .

This map is injective. Indeed, if $C, C' \in \widehat{T}_0$ admit the same supremum s in T , they are both subsets of the chain $s \downarrow \cap T_0$. But full down-chains contained in a same chain in T_0 coincide when they have the same supremum in T . Finally, it is straightforward that the order induced from T under this map coincides with the order on T_0 . \square

In the next statement, the isomorphism $T \cong \widehat{T}$ is implicitly understood.

Proposition 5.15. *Let (T, \leq) be a semi-linear order such that every non-empty chain in T has a supremum and an infimum in T . Let further $T_0 \subseteq T$ be a subset such that (T_0, \leq) is semi-linear.*

If T_0 is order-dense in T , then the canonical map $\widehat{T}_0 \rightarrow T$ is an isomorphism.

Proof. We know already that there is a canonical inclusion of semi-linear orders $\widehat{T}_0 \rightarrow T$ defined by mapping $C \in \widehat{T}_0$ to its supremum in T . We need to prove that it is surjective. Choose thus some $t \in T$. We define $C = t \downarrow \cap T_0$ and observe that the supremum of C in T is t by order-density. It suffices therefore to show that C is a full down-chain in T_0 ; the fact that it is a down-chain is immediate. Suppose therefore that C admits a supremum s in T_0 . Since s is an upper bound for C , it is also an upper bound for the chain $t \downarrow$ by order-density. Thus $s \geq t$. Applying again order-density, we deduce $s = t$. It follows $s \in C$ as desired. \square

Contrary to linear orders, partial orders admit a number of unrelated natural topologies, see e.g. [Red76, §3]. We introduce a topology on an arbitrary semi-linear order (T, \leq) as follows.

Definition 5.16. Let (T, \leq) be a semi-linear order. We endow T with the topology generated by all sets $T \setminus x \uparrow$ and $x \uparrow \setminus \{x\}$, where x ranges over T .

We recall that an ordered set is a \wedge -semi-lattice if any pair $\{p, q\}$ of elements admits an infimum, denoted by $p \wedge q$. This is of course the case if the order is \wedge -complete.

We shall say that two points x, y of a topological space are *separated* by a point t if the complement of t can be partitioned into two open sets, each containing one of x, y .

Lemma 5.17. *Let (T, \leq) be a dense semi-linear order which is a \wedge -semi-lattice. Then any two distinct points can be separated by a point; in particular, T is Hausdorff.*

Proof. Let x, y be distinct points in T ; by symmetry, we can assume $x \not\leq y$ without loss of generality. Thus $x \wedge y < x$ and by order-density there is t with $x \wedge y < t < x$. Notice that $t \not\leq y$. Now t separates x and y because $x \in t \uparrow \setminus \{t\}$ and $y \in T \setminus t \uparrow$. \square

Lemma 5.18. *Let (T, \leq) be a semi-linear order which is a \wedge -semi-lattice. If T contains a countable order-dense subset, then T is second countable as a topological space.*

Proof. Let $T_0 \subseteq T$ be a countable order-dense subset. It suffices to verify that the sets $T \setminus t \uparrow$ and $t \uparrow \setminus \{t\}$ form a sub-base of the topology when t ranges over T_0 . Consider first a point y of $T \setminus x \uparrow$ for an arbitrary $x \in T$. Then $x \wedge y < x$ and therefore there is $t \in T_0$ with $x \wedge y < t < x$. In particular, $y \in T \setminus t \uparrow$ and $T \setminus t \uparrow \subseteq T \setminus x \uparrow$. Next, consider $y \in x \uparrow \setminus \{x\}$. There is now $t \in T_0$ with $x < t < y$ and $y \in t \uparrow \setminus \{t\} \subseteq x \uparrow \setminus \{x\}$. \square

We shall use again the more precise fact established above that the sets $T \setminus t \uparrow$ and $t \uparrow \setminus \{t\}$ form a sub-base when t ranges over T_0 .

Theorem 5.19. *Let (T, \leq) be semi-linear order admitting a countable order-dense subset. Suppose that every non-empty chain in T has a supremum in T and that T has a minimum.*

Then T is a dendrite for the topology that we introduced.

Proof. By Remark 5.9, $T \cong \widehat{T}$. Therefore Proposition 5.7 implies that (T, \leq) is \wedge -complete. In particular, T is Hausdorff and second countable by Lemma 5.17 and Lemma 5.18.

Moreover, every maximal chain $\mathcal{C} \subseteq T$ is Dedekind-complete by Proposition 5.7; since moreover \mathcal{C} admits a countable order-dense subset and has a maximum and a minimum, it is order-isomorphic to a closed interval by Cantor's theorem [Can95, p. 511]. Our topology is the usual order-topology when restricted to \mathcal{C} and it follows that every point of T can be connected to the minimum by an arc.

We claim that T is compact. Being second countable, it suffices to find an accumulation point for an arbitrary sequence (x_n) in T . Suppose for a contradiction that there is no accumulation point and let $T_0 \subseteq T$ be a countable order-dense subset. Upon extracting a subsequence for each $t \in T_0$ in a diagonal process, we can assume that (x_n) has the following property. For every $t \in T_0$ there is an integer n_t such that one of the following holds: either $\forall n \geq n_t : x_n > t$, or $\forall n \geq n_t : x_n \not\leq t$.

Let $C \subseteq T$ be the collection of all $x \in T$ such that $x_n > x$ holds for all but finitely many n ; in particular C contains all $t \in T_0$ satisfying the first case of the above alternative. We can assume $C \neq \emptyset$ because if C does not contain the minimum z of T then $x_n = z$ for n large enough. The set C is a chain since any two of its elements belong to some $x_n \downarrow$. Therefore, C admits a supremum s in T and we proceed to show that x_n converges to s using the sub-base of neighbourhoods determined by T_0 .

To this end, consider first $t \in T_0$ such that $s \in T \setminus t \uparrow$. If x_n did not belong to this neighbourhood $T \setminus t \uparrow$ for a cofinal set of integers n , then $\forall n \geq n_t : x_n > t$ by the above alternative. This would imply $t \in C$ and thus $s \geq t$, which is absurd. Consider next $t \in T_0$ such that

$s \in t \uparrow \setminus \{t\}$. Suppose again that x_n is not almost always in $t \uparrow \setminus \{t\}$; this time, it follows that $\forall n \geq n_t : x_n \not\geq t$. On the other hand, t is not an upper bound for C ; since $s \downarrow$ is a chain, this implies that there is $x \in C$ with $x > t$. For n large enough, we have $x_n > x > t$, which is absurd.

This completes the proof that T is compact.

At this point, we know in particular that T is connected, compact, Hausdorff and second countable; thus T is a continuum. For continua, one of the equivalent characterisations of dendrites is that any two distinct points can be separated by a point, see [Nad92, 10.2]. In our case, this criterion is satisfied by Lemma 5.17. \square

If we start with a dendrite X and order as in Example 5.2 by choosing $z \in \text{Ends}(X)$, then the topology that we defined on this ordered set is the original dendrite topology by construction. It follows by naturality of the construction that every automorphism of the ordered set (X, \leq) preserves this topology. Conversely, every homeomorphism of X fixing z preserves the order. We record this as follows.

Proposition 5.20. *There is a canonical identification $\text{Homeo}_z(X) \cong \text{Aut}(X, \leq)$.* \square

We shall be more interested in certain countable subsets of X viewed as ordered sets.

Corollary 5.21. *Let X be a dendrite and $z \in \text{Ends}(X)$. Let $T \subseteq X$ be an arcwise dense subset that is invariant under the stabiliser $\text{Homeo}_z(X)$ of z and endow T with the semi-linear order determined by z . Then the natural map*

$$\text{Homeo}_z(X) \longrightarrow \text{Aut}(T, \leq)$$

is an isomorphism of groups.

Proof. The above map is a group homomorphism which is injective because T is dense in X ; we need to prove that it is onto. By naturality of the construction of \hat{T} and of the topology, every automorphism of T extends to a homeomorphism of \hat{T} . By Proposition 5.15, we have a canonical homeomorphism $\hat{T} \cong X$ and the statement follows. \square

6. GENERALIZED WAŻEWSKI DENDRITES

This section investigates homogeneity properties of the generalized Ważewski dendrites D_S . We start with a general definition.

To any finite subset F of a dendrite X we associate a finite vertex-labelled simplicial tree $\langle F \rangle$ as follows. The sub-dendrite $[F]$ is a finite tree in the topological sense, i.e. the topological realization of a finite simplicial tree. Such a simplicial tree is not unique because degree-two vertices can be added or removed without changing the topological realization. We choose for $\langle F \rangle$ to retain precisely one degree-two vertex for each element of F which is a regular point of the dendrite $[F]$. Thus, $\langle F \rangle$ is a tree whose vertex set contains F . Finally, we label the vertices of $\langle F \rangle$ by assigning to each vertex its order in X .

We observe that this is a labelling by elements of $\mathbf{N} \cup \{\infty\}$ and that the labellings that can arise in this way are precisely those which are bounded below by the degree of x in $\langle F \rangle$, i.e. by the order of x in $[F]$.

Proposition 6.1. *Fix $S \subseteq \{3, 4, \dots, \infty\}$ non-empty. Given two finite subsets $F, F' \subseteq D_S$, any isomorphism of labelled graphs $\langle F \rangle \rightarrow \langle F' \rangle$ can be extended to a homeomorphism of D_S .*

Proof. To produce a homeomorphism h of an arbitrary topological space X , it suffices to cover X by a finite family \mathcal{Y} of closed subsets and to specify for each $Y \in \mathcal{Y}$ a homeomorphism $h_Y: Y \cong h_Y(Y) \subseteq X$ such that $\{h_Y(Y)\}_{Y \in \mathcal{Y}}$ is also a cover by closed sets and with the compatibility conditions

$$h_Y|_{Y \cap Z} = h_Z|_{Y \cap Z}, \quad h_Y(Y \cap Z) = h_Y(Y) \cap h_Z(Z)$$

for all $Y, Z \in \mathcal{Y}$. In the case at hand, we use $\langle F \rangle$ to decompose D_S into a family \mathcal{Y} of sub-dendrites, as follows. First, for any adjacent vertices x, y of $\langle F \rangle$, we define $D_S(x, y)$ as the closure of the component of $D_S \setminus \{x, y\}$ containing the interior of the arc $[x, y]$. Next, for each individual vertex x of $\langle F \rangle$, we let Y_x be the union of $\{x\}$ and of all components of $D_S \setminus \{x\}$ that do not meet any of the $D_S(x, y)$ where y ranges over the vertices adjacent to x . We take for \mathcal{Y} the collection of all these $D_S(x, y)$ and Y_x .

We define likewise the decomposition \mathcal{Y}' associated to $\langle F' \rangle$ and observe that any isomorphism $\langle F \rangle \rightarrow \langle F' \rangle$ of labelled graphs induces a bijection $\mathcal{H}: \mathcal{Y} \rightarrow \mathcal{Y}'$. Moreover, the intersection of two distinct elements of \mathcal{Y} is either empty or reduces to a single point (which is a vertex of $\langle F \rangle$). Therefore, it suffices to exhibit for each $Y \in \mathcal{Y}$ a homeomorphism $h_Y: Y \rightarrow \mathcal{H}(Y)$ which has the prescribed behaviour on the intersection of Y and the vertex set of $\langle F \rangle$.

At this point, we can conclude by the well-known homogeneity properties of D_S as follows. All $D_S(x, y)$ are homeomorphic to D_S itself, and the homeomorphism can be chosen to send x and y to any given pair of distinct ends of D_S . This follows from Theorem 6.2 in [CD94], compare also Corollary 3.3 in [Cha95]. As for Y_x , it can be written as the union of $D_S(x, x')$ where we choose some $x' \in \text{Ends}(D_S)$ in each component of $D_S \setminus \{x\}$ meeting Y_x . The (possibly infinite) number of these components is the difference between the orders of x in D_S and in $[F]$, and therefore it is the same number as for $\mathcal{H}(Y_x)$. Hence the required homeomorphism $Y_x \cong \mathcal{H}(Y_x)$ is obtained by patching together the various homeomorphisms on each $D_S(x, x')$. The possibility of infinitely many such components does not raise any continuity issue since they will then form a null-sequence, just as in the proof of Lemma 2.9. \square

The following very special case of Proposition 6.1 in fact hardly different from the ingredient from [CD94] that we used in the proof, but we isolate it for further reference:

Corollary 6.2. *For any $S \subseteq \{3, 4, \dots, \infty\}$ and any $n \in S$, the action of $\text{Homeo}(D_S)$ on $\text{Br}_n(D_S)$, on $\text{Ends}(X)$ and on $\text{Reg}(X)$ is doubly transitive.* \square

An elementary consequence of double transitivity is the following.

Corollary 6.3. *Let $S \subseteq \{3, 4, \dots, \infty\}$ be a non-empty set and $n \in S \cup \{1, 2\}$. Then any element of $\text{Homeo}(D_S)$ is the product of two elements fixing each some point of order n in D_S .*

This clarifies the simplicity statement of Corollary 4.5 above:

Corollary 6.4. *$\text{Homeo}(D_S)$ is a simple group for any non-empty $S \subseteq \{3, 4, \dots, \infty\}$.*

Proof. By Corollary 4.5, G_{Br} is a simple group, where $G = \text{Homeo}(D_S)$. On the other hand, Corollary 6.3 implies, in particular, that $G = G_{\text{Br}}$. \square

Recalling the basic fact that doubly transitive actions are primitive, we also deduce:

Corollary 6.5. *The stabiliser in $\text{Homeo}(D_S)$ of any point of D_S is maximal as a proper subgroup of $\text{Homeo}(D_S)$.* \square

Recalling that the set of branch points is always countable whilst both $\text{Reg}(D_S)$ and $\text{Ends}(D_S)$ are uncountable, Corollary 6.2 implies also the following.

Corollary 6.6. *The stabilizer of $x \in D_S$ has countable index in $\text{Homeo}(D_S)$ if and only if x is a branch point.* \square

In the case where S is finite, there are only finitely many labelled trees as above for any given number of vertices. Therefore, we deduce the following from Proposition 6.1.

Corollary 6.7. *If S is finite, then the action of $\text{Homeo}(D_S)$ on D_S (viewed simply as a set) is oligomorphic.* \square

Here *oligomorphic* means that for each $p \in \mathbf{N}$ the diagonal action of $\text{Homeo}(D_S)$ on $(D_S)^p$ has finitely many orbits [Cam90]. When S is infinite, we recover likewise from Proposition 6.1 the weaker fact that $\text{Homeo}(D_S)$ has countably many orbits on $(D_S)^p$, first established in [Cam11].

We recall that a topological group is *Roelcke pre-compact* if for every identity neighbourhood U there is a finite set F in G with $G = UFU$. This holds in particular for groups that can be represented as closed oligomorphic permutation groups of countable sets (see e.g. [ET16, §1.2]). Therefore, considering the representation of $\text{Homeo}(D_S)$ into $\text{Sym}(\text{Br}(X))$ as in Proposition 2.4, we have:

Corollary 6.8. *If S is finite, then the Polish group $\text{Homeo}(D_S)$ is Roelcke pre-compact.* \square

By a result of Evans–Tsankov, we can deduce that $\text{Homeo}(D_S)$ has the strong Kazhdan property (T) as a topological group (which is not the case for all oligomorphic groups, see [Tsa12, §6]).

Corollary 6.9. *If S is finite, then the Polish group $\text{Homeo}(D_S)$ has the strong Kazhdan property (T) as a topological group.*

Proof. In view of Theorem 1.1 in [ET16] and of the oligomorphic presentation of $\text{Homeo}(D_S)$ into $\text{Sym}(\text{Br}(X))$, it remains only to justify that $\text{Homeo}(D_S)$ has no open subgroup of finite index. This follows from the abstract simplicity of $\text{Homeo}(D_S)$ established in Corollary 6.4. \square

Another application of Proposition 6.1 provides a link with semi-linear orders:

Corollary 6.10. *Fix $S \subseteq \{3, 4, \dots, \infty\}$, $n \in S$ and any end z of D_S . Then the action of the stabiliser $\text{Homeo}_z(D_S)$ on the subset of pairs*

$$\{(x, y) : x, y \in \text{Br}_n(D_S), x \neq y \text{ and } x \in [y, z]\}$$

is transitive.

In particular, the semi-linear order (T, \leq) induced on $T = \text{Br}_n(D_S)$ as in Example 5.2 is weakly two-transitive.

We recall here that a semi-linear order (T, \leq) is called *weakly two-transitive* if $\text{Aut}(T, \leq)$ acts transitively on the set of pairs (x, y) satisfying $x < y$.

Proof of Corollary 6.10. For any such pair (x, y) , the labelled tree $\langle \{z, x, y\} \rangle$ has always the same isomorphism type: namely, the vertices z, x, y are aligned and in this order, with labels respectively $1, n, n$. Therefore the statement follows from Proposition 6.1. \square

A group G is said to have *property (OB)* if every isometric G -action on any metric space has bounded orbits. Amongst equivalent definitions is that every left-invariant metric on G is bounded, see e.g. [Ros09, 1.2]. The above results allow us to leverage a theorem from [DT09] on semi-linear orders and deduce:

Theorem 6.11. *Let $S \subseteq \{3, 4, \dots, \infty\}$ be a non-empty set and pick $z \in \text{Ends}(D_S)$. Then the group $\text{Homeo}_z(D_S)$ has property (OB).*

Proof. Choose any $n \in S$ and let $T = \text{Br}_n(D_S)$. We endow D_S and T with the semi-linear order determined by z as in Example 5.2. By Corollary 6.10, the order (T, \leq) is weakly two-transitive. By Theorem 3.1 in [DT09], it follows that the group $\text{Aut}(T, \leq)$ has property (OB). Finally, Corollary 5.21, states that $\text{Homeo}_z(D_S)$ is isomorphic to $\text{Aut}(T, \leq)$. \square

Corollary 6.12. *For any non-empty $S \subseteq \{3, 4, \dots, \infty\}$, the group $\text{Homeo}(D_S)$ has property (OB).*

Proof. Let z be an end of D_S . By Theorem 6.11, $H = \text{Homeo}_z(D_S)$ has property (OB). By Corollary 6.2 the action of $G = \text{Homeo}(D_S)$ on G/H is doubly transitive. Now Lemma 6.13 below completes the proof. \square

Lemma 6.13. *Let G be a group and $H < G$ a subgroup such that the G -action on G/H is doubly transitive. If H has property (OB), then so does G .*

The same holds more generally if H has the relative property (OB) in G , as defined in [Ros14].

Proof. The assumption on G/H is equivalent to the fact that we have $G = H \cup HgH$ for some $g \in G$; hence the statement follows from the characterisation in terms of left-invariant metrics. \square

Another application of the isomorphism afforded by Corollary 5.21 between end stabilisers in $\text{Homeo}(D_S)$ and $\text{Aut}(T, \leq)$ will be useful in Section 7, namely:

Proposition 6.14. *Let $S \subseteq \{3, 4, \dots, \infty\}$ be non-empty and let $z \in \text{Ends}(D_S)$. Let $M \triangleleft \text{Homeo}_z(D_S)$ be the normal subgroup of elements fixing pointwise some non-trivial arc containing z .*

Then M contains all proper normal subgroups of $\text{Homeo}_z(D_S)$.

Furthermore, $\text{Homeo}_z(D_S)$ has no proper subgroup of finite index.

Proof. As before we fix $n \in S$ and consider the countable semi-linear order (T, \leq) for $T = \text{Br}_n(D_S)$ as in Example 5.2, recalling that it is weakly two-transitive by Corollary 6.10. Moreover, $H = \text{Homeo}_z(D_S)$ is canonically isomorphic to $\text{Aut}(T, \leq)$ by Corollary 5.21. Therefore, we can apply Theorem 1.3 in [DHM89] which holds for all countable weakly two-transitive semi-linear orders. This result implies that every proper normal subgroup $N \triangleleft \text{Aut}(T)$ is contained in the normal subgroup $R(T) \triangleleft \text{Aut}(T)$ of elements that fix pointwise $\{y : y < t\}$ for some $t \in T$. As a subgroup of $\text{Homeo}(D_S)$, this is precisely the group M .

For the additional statement, it suffices to prove that the group H/M is infinite since every finite index subgroup contains a normal finite index subgroup. Let $g \in \text{Homeo}(D_S)$ be a homeomorphism admitting an austro-boreal arc ending at z ; in particular, $g \in H$. Such an element g exists by Lemma 6.15 below. We claim that $g \notin M$. This claim then also holds for any non-trivial power of g and hence the group generated by the image of g in H/M is indeed infinite.

To prove the claim, it suffices to observe that any arc terminating at z must meet I at more than just z because z is an end; therefore the claim follows since the only g -fixed points in I are the two extremities of I . \square

Lemma 6.15. *Any non-trivial arc of D_S is austro-boreal for some element of $\text{Homeo}(D_S)$.*

Proof. Let $[x, y]$ be a non-trivial arc of D_S . By Theorem 10.5 in [DM16], we know that there is an element g' of $\text{Homeo}(D_S)$ admitting some austro-boreal arc $[x', y']$. By [CD94, 6.2], both $D_S(x, y)$ and $D_S(x', y')$ are homeomorphic to D_S and hence to each other (these sub-dendrites were defined in the proof of Proposition 6.1). We can now define $g \in \text{Homeo}(D_S)$ by transporting the g' -action from $D_S(x', y')$ to $D_S(x, y)$ and letting g acts trivially outside $D_S(x, y)$. \square

7. NON-ISOMORPHIC HOMEOMORPHISM GROUPS

Consider any non-empty set $S \subseteq \{3, 4, \dots, \infty\}$. The initial goal of this section is to prove that the homeomorphism group of the generalised Wazewski dendrite D_S , as an abstract group, determines the set S . The proof will show that the space D_S can be recovered from the mutual positions of the stabilisers of branch points as abstract subgroups of $\text{Homeo}_x(D_S)$.

Given a point $x \in D_S$, denote by \mathcal{C}_x the set of connected components of $D_S \setminus \{x\}$. For each $C \in \mathcal{C}_x$, the closure $\bar{C} = C \cup \{x\}$ is homeomorphic to D_S by Lemma 2.14. The stabiliser $\text{Homeo}_x(D_S)$ has a natural representation to the permutation group $\text{Sym}(\mathcal{C}_x)$. Moreover, this representation is split surjective and yields a permutational wreath product:

Lemma 7.1.

$$\text{Homeo}_x(D_S) \cong \left(\prod_{C \in \mathcal{C}_x} \text{Homeo}_x(\bar{C}) \right) \rtimes \text{Sym}(\mathcal{C}_x).$$

Proof. The kernel of the representation to $\text{Sym}(\mathcal{C}_x)$ is all of $\prod_{C \in \mathcal{C}_x} \text{Homeo}_x(\bar{C})$ because of Lemma 2.9. Therefore it suffices to prove that there is a subgroup of $\text{Homeo}_x(D_S)$ mapping isomorphically onto $\text{Sym}(\mathcal{C}_x)$ under this representation.

To this end, we select an end z of D_S and choose for each $C \in \mathcal{C}$ a homeomorphism $\varphi_C: \bar{C} \rightarrow D_S$ such that $\varphi_C(x) = z$. Given a permutation $\sigma \in \text{Sym}(\mathcal{C}_x)$, we obtain $\tilde{\sigma} \in \text{Homeo}_x(D_S)$ by defining $\tilde{\sigma}$ to be $\varphi_{\sigma(C)}^{-1} \circ \varphi_C$ on \bar{C} for each $C \in \mathcal{C}_x$. The fact that $\tilde{\sigma}$ is indeed a homeomorphism even if \mathcal{C} is infinite follows from the fact that \mathcal{C} is a null-family, as in the proof of Lemma 2.9. Now $\sigma \mapsto \tilde{\sigma}$ is indeed a section of $\text{Sym}(\mathcal{C}_x)$. \square

Corollary 7.2. *If $x \in D_S$ is a point of finite order, then every finite index subgroup of $\text{Homeo}_x(D_S)$ contains the kernel of the representation onto $\text{Sym}(\mathcal{C}_x)$.*

Proof. Since $\bar{C} \cong D_S$ and since \mathcal{C}_x is finite, Proposition 6.14 implies that this kernel, namely the product $\prod_{C \in \mathcal{C}_x} \text{Homeo}_x(\bar{C})$, has no finite index (proper) subgroup. \square

Theorem 7.3. *Let $S, S' \subseteq \{3, 4, \dots, \infty\}$ be two non-empty subsets. Suppose that there is a group isomorphism*

$$\Phi: \text{Homeo}(D_S) \longrightarrow \text{Homeo}(D_{S'}).$$

Then there is a map $\varphi: D_S \rightarrow D_{S'}$ which is a Φ -equivariant homeomorphism.

In particular, Theorem 7.3 shows that S is determined by the group $\text{Homeo}(D_S)$.

Corollary 7.4. *The groups $\text{Homeo}(D_S)$ and $\text{Homeo}(D_{S'})$ are isomorphic if and only if $S = S'$. \square*

Another consequence of Theorem 7.3 is the following.

Corollary 7.5. *The group $\text{Homeo}(D_S)$ has no outer automorphisms.*

Proof of Corollary 7.5. Let Φ be an automorphism of the group $\text{Homeo}(D_S)$. Then Theorem 7.3 provides $\varphi \in \text{Homeo}(D_S)$ which is Φ -equivariant. This means by definition that for all $x \in D_S$ and all $g \in \text{Homeo}(D_S)$ we have $\varphi(gx) = \Phi(g)\varphi(x)$. In other words, $\Phi(g) = \varphi g \varphi^{-1}$; this shows that Φ is inner. \square

Proof of Theorem 7.3. Throughout the proof, we consider $\text{Homeo}(D_S)$ as acting on D_S but also on $D_{S'}$ via Φ ; we only mention Φ explicitly when a confusion could occur. We begin with the following claim:

For every branch point $x \in D_S$, the stabiliser $\text{Homeo}_x(D_S)$ fixes a branch point in $D_{S'}$; moreover this point is unique.

Indeed, consider the decomposition of $\text{Homeo}_x(D_S)$ given by Lemma 7.1 and let $n \in S$ be the order of x . Since $\bar{C} \cong D_S$, Proposition 6.14 shows that we can apply Proposition 2.12 and deduce that $\text{Homeo}_x(D_S)$ fixes a point y in $D_{S'}$. We recall that all point stabilisers are maximal by Corollary 6.5. Therefore, since Φ is an isomorphism, we deduce that the image of $\text{Homeo}_x(D_S)$ is exactly $\text{Homeo}_y(D_{S'})$. Now y must be a branch point in view of Corollary 6.6. The uniqueness follows from the maximality of stabilisers, since otherwise D_S would have two branch points y, y' with the same stabilisers; this is readily seen to contradict the double transitivity of Corollary 6.2, using that $\text{Br}_m(y)$ is arcwise dense, where m is the order of y in $D_{S'}$.

Next, we claim that the order of y is also n . It suffices to prove that n is an invariant of the abstract group $H = \text{Homeo}_x(D_S)$ amongst stabilisers of branch points. Indeed, the case $n = \infty$ is characterised as the only case when H has no index two subgroup thanks to Lemma 2.13. When $n \geq 3$ is finite, it can be recovered from the index of the minimal finite index subgroup given by Corollary 7.2.

At this point, we can already define a natural map $\varphi: \text{Br}(D_S) \rightarrow \text{Br}(D_{S'})$ that preserves the Menger–Urysohn order. This map is Φ -equivariant by construction and bijective because it is natural in Φ . We note that we have incidentally already $S = S'$. Since $\text{Br}(D_S)$ is dense in D_S , it suffices now to show that φ is uniformly continuous with respect to the uniform structure on $\text{Br}(D_S)$ induced by D_S . This then implies that φ has a continuous extension to D_S . Such a continuous extension is automatically an equivariant homeomorphism.

By Lemma 2.10, this uniform structure is generated by the entourages U_F (restricted to $\text{Br}(D_S)$), where F ranges over the finite subsets of $\text{Br}(D_S)$. Therefore, it suffices to prove that the map φ preserves the ternary relation on $\text{Br}(D_S)$ given by $x \in [y, z]$ for $x, y, z \in \text{Br}(D_S)$.

We first show that φ preserves the following weaker ternary relation: $\{x, y, z\}$ are contained in a common arc. It suffices to express this relation purely in terms the stabilizers of branch points. This can be done because $\{x, y, z\}$ are *not* contained in a common arc if and only if there is $w \neq x, y, z$ such that

$$\text{Homeo}_w(D_S) \supseteq \text{Homeo}_x(D_S) \cap \text{Homeo}_y(D_S) \cap \text{Homeo}_z(D_S).$$

The “only if” direction holds by considering the centre w of a tripod (x, y, z) . The “if” direction follows readily from the transitivity properties of Proposition 6.1.

It only remains to express the relation $x \in [y, z]$ in terms of the above weaker ternary relation on $\text{Br}(D_S)$. This is done as follows: $x \in [y, z]$ if and only if for all $w \in \text{Br}(D_S)$, either $\{w, x, y\}$ or $\{w, x, z\}$ are contained in a common arc. The “only if” direction holds by definition. For the “if” direction, suppose that $x \notin [y, z]$. Then the arc I connecting x to its first-point projection to $[y, z]$ is not reduced to a point. Therefore, we can choose

$w \in \text{Br}(D_S)$ which does not lie in any arc containing I , and thus neither $\{w, x, y\}$ nor $\{w, x, z\}$ are contained in a common arc. \square

Theorem 7.3 can also be deduced from Rubin's much more general (and correspondingly more difficult) Theorem 0.2 in [Rub89]. Indeed:

Proposition 7.6. *For any non-empty set $S \subseteq \{3, 4, \dots, \infty\}$, the pair $(D_S, \text{Homeo}(D_S))$ satisfies Rubin's condition (*).*

Recall here that a pair (X, G) , where X is a Hausdorff topological space and G is a subgroup of $\text{Homeo}(X)$, satisfies Rubin's condition (*) from [Rub89] if the following hold:

- (1) (X, G) is *regionally disrigid*: for any non-empty open subset $U \subseteq X$, there is a non-trivial $g \in G$ such that g is the identity on $X \setminus U$.
- (2) Any non-empty open subset $U \subseteq X$ contains a non-empty open subset $U_1 \subseteq U$ which is *flexible* with respect to G , i.e. for any open subsets $V, W \subseteq U_1$ with $gV \cap W \neq \emptyset$ for some $g \in G$, there is $g' \in G$ such that $g'V \cap W \neq \emptyset$ and g' is the identity on $X \setminus U$.

Proof. We claim first that every non-empty open subset $U \subseteq D_S$ contains an open subset U_1 homeomorphic to $D_S \setminus \{z\}$, where $z \in \text{Ends}(D_S)$. Indeed, by density of $\text{Ends}(D_S)$, we can choose an end $z_0 \in U$. Since z_0 is an end, it admits a neighbourhood $U_1 \subseteq U$ whose topological boundary in D_S is reduced to a single point z (compare e.g. [Nad92, 9.3]). Since the closure of U_1 is homeomorphic to D_S by Lemma 2.14, the claim follows.

It suffices to prove that the disrigidity and flexibility conditions hold for such a set U_1 . In both cases, we can work directly with the action of the stabiliser $\text{Homeo}_z(D_S)$ on D_S , then transport the resulting homeomorphisms to U_1 and extend them by the identity on $D_S \setminus U_1$. In that setting, both conditions are immediate consequences of the fact that $\text{Homeo}_z(D_S)$ acts transitively on $\text{Br}_n(D_S)$ (Proposition 6.1 or Corollary 6.10) and that $\text{Br}_n(D_S)$ is dense in D_S . \square

We expect that general dendro-minimal dendrites cannot be reconstructed from their homeomorphism groups. However, besides the case of the generalised Ważewski dendrites, this reconstruction is also possible as soon as there is a free arc:

Theorem 7.7. *Let X be a dendro-minimal dendrite with a free arc. If $G = \text{Homeo}(X)$ is isomorphic, as a group, to $G' = \text{Homeo}(X')$ for any dendro-minimal dendrite X' , then $X' \cong X$.*

Moreover, the isomorphism of groups is induced by such a homeomorphism; therefore, G has no outer automorphisms.

Notice that Theorem 7.7 does not, at first sight, fit into Rubin's setting since X and X' are not assumed to belong both to the same class (of dendro-minimal dendrites with free arcs). But in fact, the first step of the proof is to establish that the group isomorphism forces X' to admit a free arc; from then on, one can invoke Rubin's theorem.

Proof of Theorem 7.7. Let I be a maximal free arc of X . Combining the automatic continuity of [RS07] with the extreme amenability of [Pes98], it follows that any group action of $\text{Homeo}_+(I)$ on any compact metrisable space has a fixed point, see Corollary 7 in [RS07]. In particular, it follows that $\text{Homeo}_+(I)$ fixes a point in X' .

Next, we recall from Proposition 3.1 that G contains the product $\prod_{I \in \text{Free}(X)} \text{Homeo}_+(I)$ as a normal subgroup. Lemma 2.11 implies that this product group fixes a point in X' . Since it is a normal subgroup and since X' is dendro-minimal, Lemma 4.3 implies that this product

fixes $\text{Ends}(X')$ pointwise. Recalling that the set of ends is dense unless X' admits a free arc, we conclude that X' does indeed admit a free arc.

As mentioned above, this is a point where Rubin's theorem can be applied. Indeed, the condition (*) can be readily verified after observing that any non-empty open set contains some free arc, which follows from Proposition 3.2. \square

REFERENCES

- [Cam90] Peter Jephson Cameron. *Oligomorphic permutation groups*, volume 152 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1990.
- [Cam11] Riccardo Camerlo. Dendrites as Polish structures. *Proc. Amer. Math. Soc.*, 139(6):2217–2225, 2011.
- [Can95] Georg Cantor. Beiträge zur Begründung der transfiniten Mengenlehre. *Math. Ann.*, 46:481–512, 1895.
- [CD94] Włodzimierz Jan Charatonik and Anne Marie Dilks. On self-homeomorphic spaces. *Topology Appl.*, 55(3):215–238, 1994.
- [Cha95] Janusz Jerzy Charatonik. Homeomorphisms of universal dendrites. *Rend. Circ. Mat. Palermo (2)*, 44(3):457–468, 1995.
- [DHM89] Manfred Droste, Charles Wilbur Holland, Jr., and H. Dugald Macpherson. Automorphism groups of infinite semilinear orders. I. *Proc. London Math. Soc. (3)*, 58(3):454–478, 1989.
- [DM16] Bruno Duchesne and Nicolas Monod. Group actions on dendrites and curves. Preprint, [arxiv:1609.00303v2](https://arxiv.org/abs/1609.00303v2), 2016.
- [Dro85] Manfred Droste. Structure of partially ordered sets with transitive automorphism groups. *Mem. Amer. Math. Soc.*, 57(334):vi+100, 1985.
- [DT09] Manfred Droste and John Kenneth Truss. Uncountable cofinalities of automorphism groups of linear and partial orders. *Algebra Universalis*, 62(1):75–90, 2009.
- [ET16] David Mark Evans and Todor Tsankov. Free actions of free groups on countable structures and property (T). *Fund. Math.*, 232(1):49–63, 2016.
- [Kec95] Alexander Sotirios Kechris. *Classical descriptive set theory*, volume 156 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.
- [KR07] Alexander Sotirios Kechris and Christian Rosendal. Turbulence, amalgamation, and generic automorphisms of homogeneous structures. *Proc. Lond. Math. Soc. (3)*, 94(2):302–350, 2007.
- [KW30] Casimir Kuratowski and Gordon Thomas Whyburn. Sur les éléments cycliques et leurs applications. *Fund. Math.*, 16(1):305–331, 1930.
- [Mac37] Holbrook Mann MacNeille. Partially ordered sets. *Trans. Amer. Math. Soc.*, 42(3):416–460, 1937.
- [Mar89] John A. Maroli. *Tree Permutation Groups*. PhD thesis, Bowling Green State University, 1989.
- [Nad92] Sam Bernard Nadler, Jr. *Continuum theory*, volume 158 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1992. An introduction.
- [Pes98] Vladimir G. Pestov. On free actions, minimal flows, and a problem by Ellis. *Trans. Amer. Math. Soc.*, 350(10):4149–4165, 1998.
- [Red76] Robert Horace Redfield. Generalized intervals and topology. *Czechoslovak Math. J.*, 26(101)(4):527–540, 1976.
- [Ros09] Christian Rosendal. A topological version of the Bergman property. *Forum Math.*, 21(2):299–332, 2009.
- [Ros14] Christian Rosendal. Large scale geometry of metrisable groups. Preprint, [arxiv:1403.3106v1](https://arxiv.org/abs/1403.3106v1), 2014.
- [RS07] Christian Rosendal and Sławomir Solecki. Automatic continuity of homomorphisms and fixed points on metric compacta. *Israel J. Math.*, 162:349–371, 2007.
- [Rub89] Matatyahu Rubin. On the reconstruction of topological spaces from their groups of homeomorphisms. *Trans. Amer. Math. Soc.*, 312(2):487–538, 1989.
- [SU33] Otto Schreier and Stanisław Marcin Ulam. Über die Permutationsgruppe der natürlichen Zahlenfolge. *Studia Math.*, 4:134–141, 1933.
- [Tit70] Jacques Tits. Sur le groupe des automorphismes d'un arbre. In *Essays on topology and related topics (Mémoires dédiés à Georges de Rham)*, pages 188–211. Springer, New York, 1970.
- [Tsa12] Todor Tsankov. Unitary representations of oligomorphic groups. *Geom. Funct. Anal.*, 22(2):528–555, 2012.
- [Why42] Gordon Thomas Whyburn. *Analytic Topology*. American Mathematical Society Colloquium Publications, v. 28. American Mathematical Society, New York, 1942.

INSTITUT ÉLIE CARTAN, UNIVERSITÉ DE LORRAINE, NANCY, FRANCE.
E-mail address: `bruno.duchesne@univ-lorraine.fr`

EPFL, 1015 LAUSANNE, SWITZERLAND.