BOUNDED AND UNBOUNDED COHOMOLOGY OF HOMEOMORPHISM AND DIFFEOMORPHISM GROUPS

NICOLAS MONOD AND SAM NARIMAN

ABSTRACT. We determine the bounded cohomology of the group of homeomorphisms of certain low-dimensional manifolds. In particular, for the group of orientation-preserving homeomorphisms of the circle and of the closed 2-disc, it is isomorphic to the polynomial ring generated by the bounded Euler class. These seem to be the first examples of groups for which the entire bounded cohomology can be described without being trivial.

We further prove that the C^r -diffeomorphisms groups of the circle and of the closed 2-disc have the same bounded cohomology as their homeomorphism groups, so that both differ from the ordinary cohomology of C^r -diffeomorphisms when r > 1.

Finally, we determine the low-dimensional bounded cohomology of homeoand diffeomorphism of the spheres S^n and of certain 3-manifolds. In particular, we answer a question of Ghys by showing that the Euler class in $H^4(\text{Homeo}_{\circ}(S^3))$ is unbounded.

1. INTRODUCTION

Bounded cohomology H_b^{\bullet} has many motivations and applications; a few examples in geometry, topology and dynamics are [Gro82], [Ghy87], [Dup79], [EP03], [Buc07], [MS06]. If a classical invariant is bounded, then determining a representative in bounded cohomology can refine the invariant considerably, see e.g. [Ghy01], [MS04]. Contrariwise, showing that a class in not bounded is a restriction by itself, see e.g. [Lan94, Q. F.1], [Cal04].

Unfortunately, bounded cohomology remains largely elusive: as stated in [Mon06, §1], there was no group for which all of H_b^{\bullet} was known, unless it is trivial. Recent advances brought more examples where bounded cohomology is trivial and new examples where it is pathologically huge [Löh17], [FFLM21b].

Missing were examples where we can actually describe non-trivially, and understand, the entire bounded cohomology, especially for groups of relevance in geometry, topology or dynamics. *This article is a first step in that direction*.

Since our results concern groups of homeomorphisms and diffeomorphisms of manifolds, they can be compared to the very rich supply of results on the ordinary cohomology of these groups and how the latter relates to the underlying manifold. Interestingly, some of our results exhibit a close similarity to the classical case, while others stand in strong contrast. It turns out that this behaviour depends on the category: topological or smooth.

1.A. The circle S^1 . Our first results regard the circle and are much simpler to prove than for the disc; they constitute therefore an excellent warm-up for the techniques introduced in this paper.

The ordinary cohomology of the group $\text{Homeo}_{\circ}(S^1)$ of orientation-preserving homeomorphisms of the circle is known to be a polynomial ring generated by the Euler class $\mathscr{E} \in H^2$ for flat S^1 -bundles:

(1)
$$H^{\bullet}(\operatorname{Homeo}_{\circ}(S^{1})) \cong \mathbf{R}[\mathscr{E}]$$

(we take cohomology with coefficients in **R**). This can be deduced from a remarkable theorem that Thurston ([Thu74, Cor. (b) of Thm. 5]) established for any closed manifold M. It states that that the natural map $B\text{Homeo}(M) \to B\text{Homeo}^{\tau}(M)$ induces a cohomology isomorphism, wherein Homeo^{τ} denotes the *topological* group endowed with the C^0 -topology. (A group without superscript τ will always refer to the "abstract", i.e. discrete, group; see Section 2.) Thurston's theorem applies to the neutral component Homeo_{\circ} as well. In fact, we always work with Homeo_{\circ} , since this subgroup also determines the cohomology of Homeo by analysing the action of the group of components. In the case of the circle, we have $B\text{Homeo}_{\circ}^{\tau}(S^1) \simeq \mathbb{C}P^{\infty}$ and hence (1) follows.

Our first theorem is the analogue of (1) for bounded cohomology. This is probably the first case where the entire bounded cohomology of a group can be determined without being either trivial or too pathologically large to describe. For the statement, recall that \mathscr{E} is a *bounded class* by the Milnor–Wood inequality ([Mil58], [Woo71]) and that it admits a *unique* bounded representative [MM85, Cor. 2.11] in H_b^2 , the *bounded Euler class* $\mathscr{E}_b \in H_b^2$, because Homeo_o(S^1) is uniformly perfect (see [MM85, Cor. 2.11] and [EHN81, Thm. 2.3]).

Theorem 1.1. We have $H_b^{\bullet}(\text{Homeo}_{\circ}(S^1)) \cong \mathbf{R}[\mathscr{E}_b]$, the polynomial ring generated by the bounded Euler class.

Thus the comparison map $H_b^{\bullet}(\operatorname{Homeo}_{\circ}(S^1)) \to H^{\bullet}(\operatorname{Homeo}_{\circ}(S^1))$ is an isomorphism in every degree. Such a statement completely fails already for a surface Σ_g of genus g > 0. Indeed, a recent result of Bowden-Hensel-Webb ([BHW21]) shows that $H_b^2(\operatorname{Homeo}_{\circ}(\Sigma_g))$ is infinite-dimensional; by contrast, the the results of Hamstrom [Ham74] and Thurston [Thu74] imply that $H^2(\operatorname{Homeo}_{\circ}(\Sigma_g))$ vanishes. Quasimorphisms and low dimensional bounded cohomology of volume preserving diffeomorphisms and symplectomorphisms have also been extensively studied (see [GG04, Pol06, BM21] and references therein). Another example to which we shall return is a theorem due to Mann [Man20] stating that the map $H_b^2(\operatorname{Homeo}(M)) \to H^2(\operatorname{Homeo}(M))$ has a nontrivial cokernel for certain Seifert fibered 3-manifolds M.

For diffeomorphism groups, the usual group cohomology is related to deep open questions in foliation theory, in particular to the homotopy type of the Haefliger space (see [Tsu89a] and references therein). Let $\overline{BDiff}^r(M)$ be the homotopy fiber of the map

$$\eta: BDiff^{r}(M) \to BDiff^{r,\tau}(M)$$

that is induced by the identity homomorphism. For C^1 -diffeomorphisms, remarkably Tsuboi proved (see [Tsu89b]) that η is a homology isomorphism. In all regularities, Mather–Thurston's theorem ([Thu74, Thm. 5], [Mat11]) says that the space $\overline{BDiff}^r(M)$ is homology isomorphic to the space of sections of a certain bundle over M. For C^r -diffeomorphisms when r > 1, as a consequence of Mather–Thurston's theorem and a conjecture about the connectivity of the Haefliger space ([Thu74, Conjecture], [Hae10, Section 6]), it is expected that η induces a homology isomorphism for degrees less than or equal to dim(M). For regularities r > 1 and a compact manifold M, it is known that there are non-trivial Godbillon–Vey classes in $H^{\dim(M)+1}(\text{Diff}^r(M); \mathbf{R})$ (see [Bot78, Miz17]) and in particular for the case of the circle, Thurston ([Thu72]) proved that there is a surjective map

$H_2(\operatorname{Diff}^r_{\circ}(S^1); \mathbf{Z}) \twoheadrightarrow \mathbf{R} \oplus \mathbf{Z},$

where the map to \mathbf{Z} summand is induced by the Euler class and the map to \mathbf{R} is induced by the Godbillon–Vey class. It is conjectured ([Mor17]) that this map is

in fact injective. Morita ([Mor84, Mor85, Nar16]) generalised Thurston's theorem and showed that there exists a surjective map

$$H_{2n}(\operatorname{Diff}_{\circ}^{r}(S^{1}); \mathbf{Z}) \twoheadrightarrow \mathbf{R} \oplus \mathbf{Z},$$

for all n where the map to \mathbb{Z} summand is induced by the power of the Euler class. Although determining the group cohomology of $\operatorname{Diff}_{\circ}^{r}(S^{1})$ for r > 1 seems to be a very difficult open problem, the smooth group cohomology of the infinite dimensional Lie group $\operatorname{Diff}_{\circ}^{r}(S^{1})$ satisfies the van Est isomorphism ([Hae79, Page 44]) and is completely determined as

$$H^{\bullet}_{sm}(\operatorname{Diff}^{r}_{\circ}(S^{1})) \cong \mathbf{R}[\mathscr{E},\mathscr{GV}]/(\mathscr{E}\cdot\mathscr{GV}=0).$$

Similarly, the group cohomology of the piecewise linear homomeomorphisms $PL_{\circ}(S^1)$ is much richer than the group cohomology of homeomorphisms of the circle. In fact, there is a discrete analogue of Godbillon–Vey classes ([GS87]) that is non-trivial in $H^2(PL_{\circ}(S^1))$. Combining Mather–Thurston's theorem for PL foliations and the work of Peter Greenberg ([Gre87, Cor. 1.12]), one can see that $H^{\bullet}(PL_{\circ}(S^1))$ is much bigger than the ring that is generated by the Euler class.

In contrast to this classical picture, we prove that bounded cohomology does not distinguish between homeomorphisms, PL homeomorphisms and diffeomorphisms of S^1 .

Theorem 1.2. For every $r \in \mathbf{N} \cup \{\infty\}$, the inclusion $\text{Diff}^r \to \text{Homeo}$ induces an isomorphism in bounded cohomology:

$$H_b^{\bullet}(\operatorname{Diff}^r_{\circ}(S^1)) \cong H_b^{\bullet}(\operatorname{Homeo}_{\circ}(S^1)) \cong \mathbf{R}[\mathscr{E}_b].$$

The same statement holds for the piecewise-linear category:

$$H_b^{\bullet}(\mathrm{PL}_{\circ}(S^1)) \cong H_b^{\bullet}(\mathrm{Homeo}_{\circ}(S^1)) \cong \mathbf{R}[\mathscr{E}_b].$$

As we shall see, the proof itself is rather robust with respect to changes of categories of circle transformations. For instance, it also holds for the *countable* subgroup of $PL_{\circ}(S^1)$ known as Thompson's group T. Therefore, the bounded cohomology of T is also $\mathbf{R}[\mathscr{E}_b]$; this contrasts with the ordinary cohomology determined by Ghys–Sergiescu [GS87] and provides a concrete example of a countable group whose bounded cohomology is entirely known. We note that the bounded cohomology of T can also be determined by combining [Mon22] with [FFLM21a, §6].

1.B. The closed disc D^2 . Some methods introduced for the proof of Theorem 1.1 find their full use in higher dimensions. A rather more involved implementation of our strategy allows us to determine completely the bounded cohomology of Homeo_o (D^2) .

Theorem 1.3. The restriction map $\operatorname{Homeo}_{\circ}(D^2) \to \operatorname{Homeo}_{\circ}(S^1)$ induces an isomorphism in bounded cohomology.

In particular, the algebra $H_b^{\bullet}(\text{Homeo}_{\circ}(D^2))$ is isomorphic to $\mathbf{R}[\mathscr{E}_b]$.

This recalls McDuff's theorem ([McD80, Cor. 2.13]) stating that the restriction homomorphism $\operatorname{Homeo}_{\circ}(D^n) \to \operatorname{Homeo}_{\circ}(S^{n-1})$ induces an isomorphism in ordinary cohomology for all n.

It turns out that our proof is again robust when changing to the smooth category:

Theorem 1.4. The statement of Theorem 1.3 also holds for C^r diffeomorphisms, $r \in \mathbf{N} \cup \{\infty\}$.

In particular, the algebra $H_b^{\bullet}(\operatorname{Diff}^r_{\circ}(D^2))$ is also isomorphic to $\mathbf{R}[\mathscr{E}_b]$.

Bowden showed ([Bow11, Prop. 5.1]) that in ordinary group cohomology not only the Euler class but also the Godbillon–Vey class pulls back nontrivially to the second group cohomology $H^2(\text{Diff}^r_{\circ}(D^2); \mathbf{R})$ for r > 1. Hence, the comparison map $H^2_b(\text{Diff}^r_{\circ}(D^2)) \to H^2(\text{Diff}^r_{\circ}(D^2))$ is not an isomorphism for r > 1. Shigeyuki Morita mentioned to the authors the interesting fact that the Bott vanishing theorem ([Bot70]), however, implies that \mathscr{E}^4 in $H^8(\text{Diff}^r_{\circ}(D^2))$ vanishes for r > 1.

Remark 1.5. Let $\operatorname{Diff}^{r,vol}(D^2)$ be the volume preserving C^r -diffeomorphisms of D^2 for the standard volume form. It is known ([Tsu00, Cor. 3.2]) that for the restriction map $\operatorname{Diff}^{r,vol}(D^2) \to \operatorname{Diff}^{r}_{\circ}(S^1)$, the Euler class pulls back *trivially* to $H^2(\operatorname{Diff}^{r,vol}(D^2))$.

Our general strategy is to construct a semi-simplicial set X_{\bullet} , depending on the regularity r, on which $G = \text{Diff}_{\circ}^{r}(D^{2})$ acts in such a way that we decompose D^{2} into pieces, and each piece will have a stabiliser in G that is easier to handle cohomologically. We then resolve $H_{b}^{\bullet}(G)$ in terms of $H_{b}^{\bullet}(X_{\bullet})$, of $H_{b}^{\bullet}(X_{\bullet}/G)$ and of the stabilisers. In the case of D^{2} , the set X_{\bullet} consists of configurations of neighbourhood germs of chords in the disc. The complexity of these configurations, specifically the infinite possibilities for mutual intersections of chords, has limited us to n = 2.

For this strategy to actually simplify the problem, we need to show that the stabilisers are *boundedly acyclic*, meaning that their bounded cohomology vanishes in all positive degrees. We shall establish several such general bounded acyclicity results, both as tools for the above and for the case of S^n below. The first non-trivial bounded acyclicity statement is the theorem of Matsumoto–Morita [MM85] which states that the group $\text{Homeo}_c(\mathbf{R}^n)$ of *compactly supported* homeomorphisms is boundedly acyclic. That result was a refinement of Mather's acyclicity in ordinary cohomology [Mat71]. A tool for our study of homeomorphisms and diffeomorphisms of S^n is the following generalisation of the Matsumoto–Morita theorem to the case of certain manifolds that can be "displaced", and also to higher regularities.

Theorem 1.6. Let M be any closed C^r -manifold and $n \ge 1$ and let Z be C^r diffeomorphic to $M \times \mathbf{R}^n$. Then the groups $\operatorname{Homeo}_c(Z)$, $\operatorname{Homeo}_{c,\circ}(Z)$, $\operatorname{Diff}_c^r(Z)$ and $\operatorname{Diff}_{c,\circ}^r(Z)$ are boundedly acyclic for all $r \in \mathbf{N} \cup \{\infty\}$.

We note that earlier instances of the "displacement" technique, in degree two, include Kotschick [Kot08] and Burago–Ivanov–Polterovich [BIP08] (notably their notion of portable manifold).

1.C. Flat S^n -bundles and Ghys's question. Recall that the Milnor–Wood inequality ([Mil58], [Woo71]) for a C^0 -flat S^1 -bundle $E \xrightarrow{p} \Sigma_g$ over a closed oriented surface Σ_q of genus g > 0 states

$$|\langle \mathscr{E}(p), [\Sigma_g] \rangle| \le 2g - 2.$$

Milnor proved this inequality when the flat circle bundle is linear meaning that the monodromy group of the bundle lies in $PSL_2(\mathbf{R})$ and Wood proved the case where the monodromy group lies in $Homeo_{\circ}(S^1)$. In the bounded cohomology language, it says that the Euler class is a bounded class and has a Gromov norm equal to $\frac{1}{2}$. Ghys [Lan94, § F.1] asked the following question about a generalisation of the Milnor–Wood inequality to flat S^3 -bundles:

Question 1.7. Let M^4 be a compact orientable 4-manifold and $\pi_1(M) \xrightarrow{\rho}$ Homeo_o (S^3) be a representation. Is it true that the Euler number of the associated S^3 -bundle over M is bounded by a number depending only on M? Is $\mathscr{E} \in H^4(\text{Homeo}_o(S^3))$ a bounded class?

One can define the Euler class for oriented S^3 -bundles using classifying spaces as follows. Note that any oriented (not necessarily flat) S^3 -bundle over the 4manifold M up to isomorphism corresponds to a map $M \to B$ Homeo_o $(S^3)^{\tau}$ which is well-defined up to homotopy. By Smale's conjecture in the topological category ([Hat83]), we know that Homeo_o $(S^3)^{\tau} \simeq SO(4)$. Hence, there is a universal Euler class \mathscr{E} in $H^4(BSO(4)) \cong H^4(B$ Homeo_o $(S^3)^{\tau}$) and in fact $H^4(BSO(4)) \cong \mathbb{R}^2$ which is generated by the universal first Pontryagin class \mathscr{P}_1 and \mathscr{E} (see also [Nar16, Remark 3.3]). Now as a consequence of Thurston's theorem ([Thu74, Cor. (b) of Thm. 5]), we know the natural map

 $H^4(B\operatorname{Homeo}_{\circ}(S^3)^{\tau}) \to H^4(\operatorname{Homeo}_{\circ}(S^3)),$

is an isomorphism. Therefore, the pull back of the universal \mathscr{P}_1 and \mathscr{E} on the universal C^0 -flat S^3 -bundle are nontrivial. By the abuse of notation, we denote the pull back of these classes to flat bundles by the same notation. So $H^4(\operatorname{Homeo}_{\circ}(S^3))$ is also generated by \mathscr{P}_1 and \mathscr{E} .

Back to Ghys's question above, proving the boundedness of \mathscr{E} is the approach suggested by the Gromov–Hirzebruch proportionality principle. This boundedness would generalize the Milnor–Wood inequality and be compatible with the Hirsch– Thurston theorem [HT75] stating that the Euler class vanishes when $\pi_1(M)$ is amenable. Moreover, for *linear* sphere bundles, the boundedness does indeed hold, as proved by Sullivan [Sul76] and Smillie [Smi] (see also [IT82] for this boundedness, and [BM12] for the exact bound).

In Section 6, we use the higher dimensional version of the semi-simplicial set we used for the circle case to answer Ghys's question in the negative:

Theorem 1.8. We have $H_b^4(\operatorname{Homeo}_{\circ}(S^3)) = 0$ and $H_b^4(\operatorname{Diff}_{\circ}^r(S^3)) = 0$ for all $r \neq 4$. In particular, the Euler class and the first Pontryagin class \mathscr{P}_1 in $H^4(\operatorname{Homeo}_{\circ}(S^3))$ are unbounded.

Remark 1.9. Our proof of the unboundedness of the Euler class for oriented C^0 flat S^3 -bundles is not constructive. In particular, it would be very interesting to construct explicit families of flat S^3 -bundles over a given 4-manifold with unbounded Euler number.

As we shall see, it is not hard to use the same semi-simplicial set to prove:

Theorem 1.10. We have $H_b^2(\text{Homeo}_{\circ}(S^n)) = 0$ and $H_b^3(\text{Homeo}_{\circ}(S^n)) = 0$ for all n > 1.

The same holds for $\operatorname{Diff}_{\circ}^{r}(S^{n})$ with $r \in \mathbb{N} \cup \{\infty\}$.

But to get the calculation up to H_b^4 , we will use the homotopy type of the group of diffeomorphisms of 3-dimensional pair of pants and also uniform perfectness of such groups in dimension 3. To approach the same calculation for S^2 , the main obstacle is that uniform perfectness is likely to fail in that context, in view of the work of Bowden–Hensel–Webb [BHW21]. For higher dimensional spheres, the main obstacle is the homotopy type of group of diffeomorphisms of higher dimensional pair of pants.

In a first draft of this paper we established Theorem 1.10 for n = 2, 3 only, due to a restriction on stabilisers appearing for the semi-simplicial sets. Thanks to a result of Fournier-Facio and Lodha [FFL21], this restriction is lifted. Likewise, we did not cover the case of diffeomorphisms. That generalisation was made possible by the recent note [Mon22] (which also lifts the restriction on n). Acknowledgements. Upon becoming aware of our work, Fournier-Facio, Löh and Moraschini have kindly shown us a draft of their work [FFLM21a]. We are grateful for their comments. Although the two lines of investigation have essentially no overlap, their work gives an alternative proof of the bounded acyclicity of some of the stabilisers appearing within some proofs below. We are grateful to Jonathan Bowden for his comments on the group of homeomorphisms and diffeomorphisms of 3-dimensional pair of pants. We thank Shigeyuki Morita for his comment about the powers of the Euler class for flat D^2 -bundles and also we thank Mehdi Yazdi for his careful reading and critical comments on the earlier draft of this paper. We are grateful to the anonymous referee for many comments which improved our exposition.

S.N. was partially supported by NSF DMS-2113828 and Simons Foundation (855209, SN).

2. NOTATION AND BACKGROUND RESULTS

We write $D^n \subseteq \mathbf{R}^n$ for the closed *n*-dimensional disc (ball), $B^n = \operatorname{int}(D^n)$ for the open one, and $S^{n-1} = \partial D^n$ for the (n-1)-sphere.

Notations such as Homeo, Diff^{*r*} and their variants shall refer to the corresponding groups, without any additional topological structure. If we drop the order of regularity *r* for diffeomorphisms, we mean smooth diffeomorphisms. If we endow them with a topology, in this case the C^0 or C^r -topology, then we shall write Homeo^{τ}, Diff^{*r*, τ} etc. for the resulting topological groups. We warn the reader that several authors adopt the opposite convention, where the groups are topologized by default and a superscript δ is added to indicate that e.g. Homeo^{δ} is stripped of its topology, or endowed with the discrete topology.

(The present convention, arguably more pedantically precise, is convenient here since the objects of study are "abstract" groups of homeomorphism; it leads occasionally to an unusual notation such as $SO(3)^{\tau}$ for the *compact* group of rotations.)

Given a subset $Y \subseteq X$ of a topological space X, we denote by Homeo(X;Y) the subgroup of Homeo(X) consisting of those homeomorphisms that fix Y pointwise. We denote by Homeo(X; near Y) the subgroup of Homeo(X;Y) of all elements that fix pointwise a neighbourhood of Y; this neighbourhood depends a priori on the element.

If no coefficients are explicitly indicated, then cohomology, bounded cohomology and any function spaces are always understood with coefficients in \mathbf{R} viewed as a trivial module.

Given a semi-simplicial set X_{\bullet} , the **bounded cohomology** $H_b^{\bullet}(X_{\bullet})$ of X_{\bullet} refers to the cohomology of the associated complex of spaces of bounded functions

$$0 \longrightarrow \ell^{\infty}(X_0) \longrightarrow \ell^{\infty}(X_1) \longrightarrow \ell^{\infty}(X_2) \longrightarrow \cdots$$

Although H_b^n is still poorly understood for general n, we note that X_{\bullet} is connected if and only if $H_b^0(X_{\bullet})$ has dimension one. The semi-simplicial set is called **boundedly** acyclic if $H_b^n(X_{\bullet})$ vanishes for all n > 0.

The bounded cohomology $H_b^{\bullet}(G)$ of a group G and the corresponding notion of bounded acyclicity are obtained by taking X_{\bullet} to be the Milnor join of G. Equivalently, the *homogeneous* resolution leads to an identification of $H_b^{\bullet}(G)$ with the cohomology of the complex of invariants

$$0 \longrightarrow \ell^{\infty}(G)^G \longrightarrow \ell^{\infty}(G^2)^G \longrightarrow \ell^{\infty}(G^3)^G \longrightarrow \cdots$$

where G acts diagonally on $G^{\bullet+1}$ and where the differentials are given by the simplicial Alexander–Kolmogorov–Spanier face maps (alternating sums omitting variables).

BOUNDED COHOMOLOGY OF HOMEOMORPHISM AND DIFFEOMORPHISM GROUPS 7

The following fundamental result of Matsumoto–Morita will be used extensively.

Theorem 2.1 (Matsumoto–Morita, Thm. 3.1 in [MM85]). The group Homeo_c(\mathbf{R}^n) of compactly supported homeomorphisms of \mathbf{R}^n (equivalently of B^n) is boundedly acyclic.

That result has recently been extended by various authors to other and wider settings. First, to all *mitotic* groups [Löh17]. Then, at the same time as we wrote the first version of the present article, to *binate* groups [FFLM21a]. After this, a criterion was established in [Mon22] which covers the following situation including numerous homeomorphism or diffeomorphism groups:

Theorem 2.2 (Cor. 6 in [Mon22]). Let G be a group acting faithfully on a set Z. Suppose that Z contains a subset Z_0 and that G contains an element $g \in G$ with the following properties:

- (i) every finite subset of G can be conjugated so that all its elements are supported in Z_0 ;
- (ii) $g^p(Z_0)$ is disjoint from Z_0 for every integer $p \ge 1$.

Then G is boundedly acyclic.

In general we have no reason to believe that an infinite product of boundedly acyclic groups is necessarily boundedly acyclic. However, the above sufficient condition is stable under products; indeed, it suffices to consider the action of the product on the corresponding disjoint union Z of sets, and to define Z_0 as the disjoint union of the corresponding subsets. Hence, we record the following.

Lemma 2.3. Any product of groups satisfying the conditions of Theorem 2.2 will satisfy them too, and hence be boundedly acyclic. \Box

Next, we state a fundamental stability result for the vanishing of bounded cohomology (without any claim of originality); it implies in particular that bounded acyclicity is preserved under finite direct products.

Proposition 2.4. Let G be a group, $K \triangleleft G$ a normal subgroup and $N \in \mathbb{N} \cup \{\infty\}$. If $H_b^n(K)$ vanishes for all 0 < n < N, then there is an isomorphism $H_b^n(G) \cong H_b^n(G/K)$ for all $0 \le n < N$.

Proof. This follows from a bounded version of the Lyndon–Hochschild–Serre spectral sequence, see e.g. [Mon01, §12]. For the reader's convenience, we expand on the necessary details:

Define Q = G/K and consider the double complex

$$\ell^{\infty}(G^{p+1} \times Q^{q+1})^{G} \cong \ell^{\infty}(G^{p+1}, \ell^{\infty}(Q^{q+1}))^{G} \cong \ell^{\infty}(Q^{q+1}, \ell^{\infty}(G^{p+1}))^{G}$$

indexed by $p, q \ge 0$, with the two differentials being given by the usual homogeneous differential on the variables in G and in Q respectively. We are now in a very special case of the setting considered in [Mon01, §12], the differences being as follows. (1) The reference allows for non-trivial coefficients F, while here $F = \mathbf{R}$. (2) The reference considers locally compact groups acting on measure spaces S, T, while here all groups are discrete and simply act on themselves, so S = G and T = Q. (3) The reference assumes that the locally compact groups are second countable in order to avoid difficulties with measurability; this assumption is not needed here since we have no measurability questions.

We can therefore quote from [Mon01, §12] as follows. The double complex gives rise to two spectral sequences, the first of which collapses already in the first page and abuts to the bounded cohomology $H_b^{\bullet}(G)$ of G, see Prop. 12.2.1

in [Mon01]. The second spectral sequence yields the same limit up to isomorphism and Prop. 12.2.2(ii) in [Mon01] states that the second page of this sequence is

$$E_2^{p,q} \cong H_b^p(Q, H_b^q(K))$$

as soon as $H_b^q(K)$ is Hausdorff. Since $H_b^0(K) \cong \mathbf{R}$ is always Hausdorff, $E_2^{p,0}$ is isomorphic to $H_b^p(Q)$ and our vanishing assumption on $H_b^n(K)$ allows us to deduce that $E_2^{p,q}$ vanishes for 0 < q < N and all p. These facts together imply the stated isomorphisms $H_b^n(G) \cong H_b^n(G/K)$ for all $0 \le n < N$.

In more elaborate vanishing arguments, it will be crucial to have a more refined, quantitative, control of the vanishing. To that end, we introduce the notion of *vanishing moduli* as follows.

Definition 2.5. Let X_{\bullet} be a semi-simplicial set satisfying $H_b^n(X_{\bullet}) = 0$ for some *n*. Then we define the *n*th vanishing modulus of X_{\bullet} as

$$\sup_{c \in C_1} \inf \left\{ \|b\|_{\infty} : b \in \ell^{\infty}(X_{n-1}) \text{ with } db = c \right\},$$

where

$$C_1 = \Big\{ c \in \ell^{\infty}(X_n) \text{ with } dc = 0 \text{ and } \|c\|_{\infty} \le 1 \Big\}.$$

In other words, the vanishing modulus controls the smallest norm of a cochain b that witnesses that a given cocycle c is a coboundary. This modulus is finite by the open mapping theorem applied to the coboundary map d. This quantity is dual to the "uniform boundary condition" constant considered by Matsumoto– Morita [MM85], although they consider it more generally for the trivial cycles within possibly non-acyclic normed complexes.

In particular, if G is a group with vanishing $H_b^n(G)$, we can speak of the *n*th vanishing modulus of G. It can be computed either on the Milnor join of G or on the homogeneous resolution since they give isometric cochain complexes.

A first occurrence of this notion is as follows. The direct factors of a boundedly acyclic product of groups are boundedly acyclic, but we will need the following uniformity statement which is a priori stronger for infinite products.

Proposition 2.6. Fix a positive integer q and let $(G_i)_{i \in I}$ be any family of groups. If the product $\prod_{i \in I} G_i$ is boundedly acyclic, then the qth vanishing modulus of all subproducts $\prod_{i \in J} G_j$ is bounded independently of $J \subseteq I$.

In particular, the qth vanishing modulus of all G_i is bounded independently of i.

The proof uses the following elementary technical observation, which we isolate for later reference.

Lemma 2.7. Let G be a group and $G_1 < G$ a subgroup such that $H_b^q(G_1) = 0$ for some q > 0. Then the qth vanishing modulus of G_1 coincides with the qth vanishing modulus of the semi-simplicial orbit set given by $(G^{p+1})/G_1$ in each dimension p.

Proof of Lemma 2.7. It is known that the bounded cohomology $H_b^{\bullet}(G_1)$ can be realised *isometrically* on the homogeneous resolution

 $\cdots \longrightarrow \ell^{\infty}(G^q)^{G_1} \longrightarrow \ell^{\infty}(G^{q+1})^{G_1} \longrightarrow \ell^{\infty}(G^{q+2})^{G_1} \longrightarrow \cdots$

see e.g. [Mon01, 7.4.10] for a more general statement. The proof given in this reference proceeds by exhibiting maps between this resolution and the homogeneous resolution for G_1 which are non-expanding already at the cocycle level, which implies the statement of the lemma.

We point out that in the case considered here, the technical tools used in [Mon01, 7.4.10] simply boil down to extending functions from G_1^{q+1} to G^{q+1} by using coset representatives.

Proof of Proposition 2.6. We claim that for a boundedly acyclic product $G = G_1 \times G_2$ of two groups, the factor G_1 is boundedly acyclic with its *q*th vanishing modulus bounded by that of *G*. This claim implies the statement of the proposition by regrouping the factors in the (possibly infinite) product.

Recall that inflation refers to the morphism $H_b^{\bullet}(G_1) \to H_b^{\bullet}(G)$ induced by the projection $G \to G_1$ while restriction is the morphism $H_b^{\bullet}(G) \to H_b^{\bullet}(G_1)$ induced by the inclusion $G_1 \to G$; thus their composition gives the identity on $H_b^{\bullet}(G_1)$. Moreover, the inclusion of G_1 -invariants into G-invariants realizes the inflation map. Therefore, we can compute the norms of the relevant cocycles and coboundaries while realising the inflation-restriction morphisms by the following two inclusions of (differential complexes of) Banach spaces:

$$\ell^{\infty}((G_1)^{q+1})^{G_1} = \ell^{\infty}((G/G_2)^{q+1})^G \subseteq \ell^{\infty}(G^{q+1})^G \subseteq \ell^{\infty}(G^{q+1})^{G_1}$$

The fact that we obtain the correct norms also in the right hand side follows from Lemma 2.7 and hence the claim is established. $\hfill\square$

3. Bounded cohomology of semi-simplicial sets

The following is a versatile method for constructing boundedly acyclic semisimplicial sets.

Definition 3.1. Let X be a set and let \perp be any binary relation on X. We say that the relation \perp is **generic** if, given any finite set $F \subseteq X$, there exists $x \in X$ with $x \perp y$ for all $y \in F$.

Furthermore, we define a semi-simplicial set X_{\bullet}^{\perp} as follows: X_{n}^{\perp} consists of all (n+1)-tuples (x_{0}, \ldots, x_{n}) in X^{n+1} such that $x_{i} \perp x_{j}$ holds for all i < j. Note that X_{n}^{\perp} is non-empty when \perp is generic (and $X \neq \emptyset$). The face maps are defined to be the usual simplex face maps.

In general \perp will be thought to represent a suitable transversality condition. A basic example is the relation \neq , which is generic as soon as X is infinite. A non-symmetric example is the relation < on $X = \mathbf{R}$. We shall later see more interesting examples with tuples of germs of arcs.

We emphasize that even though X_{\bullet}^{\perp} consists of pairwise- \perp tuples, the definition of genericity must be verified for *all* finite sets F, without assuming any relation between their elements.

Proposition 3.2. If \perp is generic, then X_{\bullet}^{\perp} is boundedly acyclic (and connected).

Proof. Given $F \subseteq X$ finite, let $V_F = \{x \in X : x \perp y \forall y \in F\}$. Note that $V_F \cap V_{F'} \supseteq V_{F \cup F'}$. Thus our assumption implies that the collection of all such V_F is a proper filter base on X. Choose an ultrafilter U on X containing the filter generated by this base. For any *bounded* function f on X, we can therefore consider the ultralimit $\lim_{x \to U} f(x)$.

We define a map

$$h_n: \ell^{\infty}(X_n^{\perp}) \longrightarrow \ell^{\infty}(X_{n-1}^{\perp})$$

by

$$(h_n f)(x_1,\ldots,x_n) = \lim_{t \to T} f(x,x_1,\ldots,x_n)$$

We note that h_n is linear and bounded (of norm one); the crucial point is that it is well-defined, since the collection of x for which $f(x, x_1, \ldots, x_n)$ is defined belongs to U. It is now routine to verify that h_{\bullet} provides a contracting homotopy, because for any given x_1, \ldots, x_n and x as above, the points x, x_1, \ldots, x_n determine a full simplex in X_{\bullet}^{\perp} . Explicitly, recall that the differential $\ell^{\infty}(X_{n-1}^{\perp}) \to \ell^{\infty}(X_n^{\perp})$ is the alternating sum of the maps $d_{n,i}$ over $0 \le i \le n$, where $d_{n,i}$ omits the *i*th element of a (n+1)-tuple. Then the relations $d_{n,i}h_n = h_{n+1}d_{n+1,i+1}$ and $h_{n+1}d_{n+1,0} = \text{Id hold}$, which shows that h_{\bullet} is a contracting homotopy.

Let G be a group acting on a semi-simplicial set X_{\bullet} . This gives rise to hypercohomology spectral sequences relating the bounded cohomology of G to that of X_{\bullet} and of its quotient X_{\bullet}/G , stated in Theorem 3.3 below. Some care is needed since a number of standard cohomological techniques do not hold for bounded cohomology. Definition 2.5 affords us a quantitative control.

Theorem 3.3. Let G be a group acting on a boundedly acyclic connected semisimplicial set X_{\bullet} and let $N \in \mathbb{N} \cup \{\infty\}$. We make the following assumptions for each $0 \le p < N$:

- (i) The stabiliser of any point in X_p has vanishing H_b^q for all q > 0 with p+q < N.
- (ii) Given q > 0 with p + q < N, the qth vanishing moduli of all those stabilisers are uniformly bounded.

Then there is an isomorphism $H_h^p(G) \cong H_h^p(X_{\bullet}/G)$ for every $0 \le p < N$.

Remark 3.4. In order to demystify the condition (ii), we note that it is obviously satisfied when for instance there are finitely many group isomorphism classes among the stabilisers in G of points in X_p . In the even more special case where there are only finitely many conjugacy classes in G of such stabilisers, the above theorem admits a simpler proof which does not explicitly involve vanishing moduli. In any case, we shall also need to apply it to situations with infinitely many isomorphism classes of stabilisers.

The technical assumption (ii) is made to ensure the uniform boundedness entering the following.

Lemma 3.5. Under the assumptions of Theorem 3.3, consider the G-representation on $\ell^{\infty}(X_p)$. Then $H^q_h(G, \ell^{\infty}(X_p))$ vanishes for all q > 0 with p + q < N.

Proof of Lemma 3.5. Fix p and q as in the statement. Let $J \subseteq X_p$ be a set of representatives of the G-orbits and let $G_j < G$ be the stabiliser of $j \in J$. Thus X_p can be identified with $\bigsqcup_{i \in J} G/G_i$.

Consider now a cocycle c representing an element $H_b^q(G, \ell^{\infty}(X_p))$; that is, c is a G-equivariant bounded cocycle $c: G^{q+1} \to \ell^{\infty}(X_p)$. The above decomposition of X_p decomposes c as into G-equivariant bounded cocycles $c_j: G^{q+1} \to \ell^{\infty}(G/G_j)$ and moreover $\|c\|_{\infty} = \sup_{j \in J} \|c_j\|_{\infty}$. In order to prove the statement, it suffices to show that each c_j is a coboundary of some bounded G-equivariant $b_j: G^q \to \ell^{\infty}(G/G_j)$ in such a way that $\|b_j\|_{\infty}$ is bounded *independently* of j; indeed in that case the map $b: G^q \to \ell^{\infty}(X_p)$ defined by these b_j will itself be bounded and witness c = db. We thus fix some $j \in J$ and proceed to show that fact:

Claim. Given c_j as above, consider the corresponding G_j -invariant map $\overline{c_j}: G^{q+1} \to \mathbf{R}$ defined by $\overline{c_j}(g_0, \ldots, g_q) = c_j(g_0, \ldots, g_q)(e)$. Then $c_j \mapsto \overline{c_j}$ is an isometric isomorphism on the cochain level from the standard resolution for $H^q_b(G, \ell^{\infty}(G/G_j))$ to the resolution for $H^q_b(G_j)$ given by bounded G_j -invariant cochains on G^{q+1} (as in Lemma 2.7).

This claim is a very explicit form of the Eckmann–Shapiro isomorphism for bounded cohomology, on the cochain level, and it is proved e.g. in [Mon01, 10.1]. With this claim in hand, the assumptions of Theorem 3.3 can now be applied to obtain $b_j: G^q \to \ell^{\infty}(G/G_j)$ with $\|b_j\|_{\infty}$ bounded independently of j, as desired. \Box

Proof of Theorem 3.3. We consider the double complex

$$L^{p,q} \coloneqq \ell^{\infty} (G^{p+1} \times X_q)^G \cong \ell^{\infty} (G^{p+1}, \ell^{\infty} (X_q))^G.$$

The spectral sequence associated to the horizontal filtration of $L^{\bullet,\bullet}$ has E_1 -page defined by the cohomology of

$$\ell^{\infty}\left(G^{p+1},\ell^{\infty}(X_{q-1})\right)^{G} \longrightarrow \ell^{\infty}\left(G^{p+1},\ell^{\infty}(X_{q})\right)^{G} \longrightarrow \ell^{\infty}\left(G^{p+1},\ell^{\infty}(X_{q+1})\right)^{G}$$

Since X_{\bullet} is boundedly acyclic, this cohomology vanishes for all $p \ge 0$ and all q > 0because for fixed p the functor $\ell^{\infty}(G^{p+1}, -)^{G}$ is exact for dual morphisms (see e.g. [Mon01, 8.2.5], wherein the second countability assumption is irrelevant in our setting since we consider G without topology). For q = 0, we obtain $\ell^{\infty}(G^{p+1})^G$ by connectedness of X_{\bullet} . It follows that the spectral sequence converges to $H_{h}^{\bullet}(G)$.

Turning to the spectral sequence associated to the vertical filtration, the E_1 -page is

$$E_1^{p,q} = H_h^q \left(G, \ell^\infty(X_p) \right)$$

 $E_1^{p,q} = H_b^q(G, \ell^{\infty}(X_p)).$ Applying Lemma 3.5 to $X = X_p$, the assumption (i) and (ii) imply $E_1^{p,q} = 0$ for all q > 0 with p + q < N. As for $E_1^{p,0}$, we have

$$E_1^{p,0} = \ell^\infty (X_p)^G \cong \ell^\infty (X_p/G)$$

which implies

$$E_2^{p,0} \cong H_b^p(X_{\bullet}/G).$$

Putting everything together, we have as claimed $H_h^p(G) \cong H_h^p(X_{\bullet}/G)$ for $0 \le p <$ N.

4. Homeomorphisms and diffeomorphisms of the circle

The case of the circle introduces already some of the ideas that will be used repeatedly in this article, but it evades technical considerations such as moduli of vanishing or difficult stabilisers. In fact, even the generic relation below is not strictly needed in this case, although it does simplify the set of stabilisers and the quotient.

We begin with a general definition: a fat point in an oriented n-manifold Mrefers to a germ at 0 of an orientation-preserving embedding $B^n \to M$, where $B^n \subseteq \mathbf{R}^n$ is the open unit disc. The image of 0 in M is the **core** of the fat point.

We shall prove Theorem 1.1 using a semi-simplicial set of fat points. Let thus X be the set of fat points in S^1 . Given $x, y \in X$, we write $x \perp y$ if x and y have distinct cores. This is a generic relation; therefore, Proposition 3.2 implies that X_{\perp}^{\perp} is a boundedly acyclic connected semi-simplicial set.

Proof of Theorem 1.1. Since the action of $G = \text{Homeo}_{\circ}(S^1)$ on X preserves the relation \perp , it induces an action on the semi-simplicial set X_{\perp}^{\perp} . We claim that the conditions of Theorem 3.3 are satisfied for $N = \infty$.

Regarding condition (i), observe that the stabiliser in G of a fat point is isomorphic to the group $Homeo_c(\mathbf{R})$ of compactly supported homeomorphisms, which is boundedly acyclic by Matsumoto–Morita's Theorem 2.1.

More generally, let L be the stabiliser of an element of X_p^{\perp} . The cores to the fat points determine p + 1 components of the circle and these components cannot be permuted by L since every element of L is trivial in some neighbourhood of the cut-points. Thus L is isomorphic to the direct product of p+1 copies of Homeo_c(**R**). Therefore, in view of Proposition 2.4, this subgroup is also boundedly acyclic.

As to condition (ii), the above discussion of stabilisers already establishes that there are finitely many isomorphism types for each given p. We observe, with Remark 3.4 in mind, that we have the stronger fact that there are only finitely many *G*-orbits in every given X_p^{\perp} . Indeed, the orbit of a tuple $(x_0, \ldots x_p)$ is determined by the cyclic ordering of the cores \dot{x}_i of the x_i . To see this, consider another tuple (y_0, \ldots, y_p) such that the \dot{y}_i have the same cyclic order as the \dot{x}_i . Choose orientation-preserving embeddings $\xi_i: B^1 \to S^1$ representing x_i in such a way that all images $\xi_i(B^1)$ are pairwise disjoint, which is possible since the \dot{x}_i are distinct. Define similarly η_i for y_i . We have partial homeomorphisms $\eta_i \circ \xi_i^{-1}$, which can be completed to a homeomorphism $h \in G$ since we have disjoint images in the same cyclic order. By construction, $h(x_i) = y_i$ holds for all i.

In conclusion, Theorem 3.3 implies that $H^{\bullet}_{\mathbf{b}}(G)$ is isomorphic to the bounded cohomology of the semi-simplicial set $X^{\perp}_{\mathbf{o}}/G$, which coincides with the usual cohomology $H^{\bullet}(X^{\perp}_{\mathbf{o}}/G)$ since all orbits sets X^{\perp}_p/G are finite. In fact we showed more, namely that $X^{\perp}_{\mathbf{o}}/G$ is the complex of cyclic orderings of finite sets, which is wellknown and easily verified to have $\mathbf{R}[\mathscr{E}]$ as its cohomology ring, see e.g. [Jek12, §5]. Alternatively, this can be seen by reversing the above argument, but in usual cohomology (see [Nar20, §3.1.3]). That is to say, $H^{\bullet}(X^{\perp}_{\mathbf{o}}/G)$ is isomorphic to $H^{\bullet}(G)$ because one checks that $X^{\perp}_{\mathbf{o}}$ is a contractible semisimplicial set, and because the stabilisers are acyclic (in the usual sense). The latter fact follows very much like in the bounded case above: it is reduced to the acyclicity of $\operatorname{Homeo}_c(\mathbf{R})$ established by Mather [Mat71] by using the Künneth formula.

Finally, the fact that $H^{\bullet}(G)$ is isomorphic to $\mathbb{R}[\mathscr{E}]$ follows from Thurston's theorem ([Thu74, Cor. (b) of Thm. 5]) as recalled in the introduction. All the above arguments being completely natural and given by a morphism of spectral sequences (the one for bounded cohomology is mapped to the ordinary one by the forgetful functor), the isomorphism preserves also the ring structure of [bounded] cohomology.

Proof of Theorem 1.2. We now turn to the C^r -diffeomorphism group $G^r = \text{Diff}^r_o(S^1)$, where $r \in \mathbf{N} \cup \{\infty\}$. In analogy with the topological case, we define C^r -fat points as germs of C^r -embeddings. This yields a corresponding semi-simplicial G^r -set $C^r X^{\perp}_{\bullet}$ which is still boundedly acyclic by Proposition 3.2. The quotient $C^r X^{\perp}_{\bullet}/G^r$ is isomorphic to the quotient X^{\perp}_{\bullet}/G of the proof of Theorem 1.1 because the same transitivity property holds, namely: the orbit of a tuple is determined by the cyclic order of the cores of the fat points constituting the tuple. A priori, the only substantial difference between the two settings resides with the stabilisers, which are products of copies of $\text{Diff}^r_c(\mathbf{R})$. Indeed, recall that $\text{Diff}^r_c(\mathbf{R})$ is not an acyclic group for r > 1 since there is the Godbillon–Vey class $\mathscr{GV} \in H^2(\text{Diff}^r_c(\mathbf{R}))$ which is nontrivial. However, this group is still boundedly acyclic by Theorem 4.2 below. Therefore we conclude as above that $H^{\bullet}_{\bullet}(G)$ is isomorphic to $\mathbf{R}[\mathscr{E}_b]$.

In order to complete the proof of Theorem 1.2 in the C^r case, is only remains to justify that the isomorphism is induced by the restriction map associated to the inclusion ι of $\operatorname{Diff}^r_{\circ}(S^1)$ into $\operatorname{Homeo}_{\circ}$. This is the case because the comparison between the two proofs goes through the ι -equivariant semi-simplicial inclusion morphism $C^r X^{\bullet}_{\bullet}$, which induces the identification between $C^r X^{\bullet}_{\bullet}/G^r$ and X^{\bullet}_{\bullet}/G .

Finally, in the PL case, we consider likewise PL fat points and only need to justify that $PL_c(\mathbf{R})$ is boundedly acyclic, which we do in Lemma 4.4 below. This completes the proof of Theorem 1.2.

(In fact the PL case is even simpler because it is equally easy to show the bounded acyclicity of $PL(\mathbf{R})$, so that fat points can be replaced by usual points in that instance.)

This proof can further be adapted to other circle transformation groups as long as the main ingredients are preserved. Specifically, suppose that a group G <Homeo_o(S^1) acts transitively on all cyclically oriented tuples of points in some Gorbit, or on some fat version thereof. If the stabiliser of any such tuple (respectively fat tuple) is boundedly acyclic, then $H^{\bullet}_{h}(G)$ is isomorphic to $\mathbf{R}[\mathscr{E}_{h}]$. We record the special case of Thompson's group T (see [CFP96] for a definition), where the bounded acyclicity of stabilisers is proved in [Mon22].

Corollary 4.1. The bounded cohomology ring $H_b^{\bullet}(T)$ of Thompson's group T is isomorphic to $\mathbf{R}[\mathcal{E}_b]$.

Other examples of groups for which this argument holds (again using [Mon22]) are piecewise-projective circle groups discussed in [Mon13].

We now justify the bounded acyclicity that was used in the proof of Theorem 1.2 for various compactly supported transformation groups of \mathbf{R} . We establish this in the following more general setting, as announced in Theorem 1.6.

Theorem 4.2. Let $n \in \mathbb{N}$ and consider $Z = \mathbb{R}^n$, or more generally $Z = M \times \mathbb{R}^n$ for any closed manifold M when n > 0.

Then the groups $G = \text{Homeo}_{c}(Z)$, $\text{Homeo}_{c,\circ}(Z)$, $\text{Diff}_{c}^{r}(Z)$ and $\text{Diff}_{c,\circ}^{r}(Z)$ are boundedly acyclic for all $r \in \mathbb{N} \cup \{\infty\}$.

Proof. In order to apply Theorem 2.2, we need to check the two conditions of that theorem. Let $Z_0 = M \times D^n$, where D^n is the closed unit ball in \mathbb{R}^n .

To verify the first condition, it suffices to show that every compact set $C \subseteq M \times \mathbb{R}^n$ can be mapped into Z_0 by some element $h \in G_o$. We can take $h = \operatorname{Id}_M \times h'$, where h'is a homothety on some large ball to ensure $h'(C) \subseteq D^n$, and then radially smoothen it to the identity away from a larger ball.

The second condition postulates the existence of g in G or G_{\circ} such that $g^{p}(Z_{0})$ is disjoint from Z_{0} for every integer $p \geq 1$. For n = 1 and M trivial, take first g_{0} to be any "bump shift", that is, a transformation which is strictly increasing on some bounded open interval I containing the interval D^{1} . Then any sufficiently high power g of g_{0} will have the required property since g_{0} is order-preserving. For n > 1, we can use the one-dimensional case in one coordinate and suitably smoothen it out to the identity along the other coordinates. In the case of $M \times \mathbb{R}^{n}$, we extend g by the identity on the M coordinate.

Since this proof reduces the statement to Theorem 2.2, we can combine it with Lemma 2.3; we record this as follows:

Corollary 4.3. The powers $\operatorname{Homeo}_{c}(\mathbf{R}^{n})^{\mathbf{N}}$, $\operatorname{Diff}_{c}^{r}(\mathbf{R}^{n})^{\mathbf{N}}$ and $\operatorname{Diff}_{c,\circ}^{r}(\mathbf{R}^{n})^{\mathbf{N}}$ are boundedly acyclic. The same holds more generally for $M \times \mathbf{R}^{n}$ instead of \mathbf{R}^{n} , where M is any closed manifold and n > 0.

If in the proof of Theorem 4.2 we replace the smooth bump shift by a PL bump shift and the (local) homothety by an analogous PL shrinking map, we obtain:

Lemma 4.4. The group $PL_c(\mathbf{R}^n)$ is boundedly acyclic for all $n \in \mathbf{N}$.

5. Homeomorphisms and diffeomorphisms of the disc

In this section, we further leverage the method of generic semi-simplicial sets to prove a bounded version in dimension n = 2 of McDuff's theorem ([McD80, Cor. 2.13]) stating that the restriction homomorphism

$$\operatorname{Homeo}_{\circ}(D^n) \to \operatorname{Homeo}_{\circ}(S^{n-1})$$

induces a cohomology isomorphism:

Theorem 5.1. The restriction map $\text{Homeo}_{\circ}(D^2) \to \text{Homeo}_{\circ}(S^1)$ induces an isomorphism in bounded cohomology.

Thus we can apply Theorem 1.1 and deduce:

Corollary 5.2. $H_b^{\bullet}(\operatorname{Homeo}_{\circ}(D^2)) \cong \mathbf{R}[\mathscr{E}_b].$

Remark 5.3. As a consequence of McDuff's result([McD80, Cor. 2.12]) at least in dimensions $n \leq 3$ we know that the inclusion Homeo_o $(D^n) \hookrightarrow$ Homeo_o (B^n) induces an isomorphism on group cohomology. Calegari ([Cal04, §4.1]) however proved that the Euler class in H^2 (Homeo_o (B^2)) is not a bounded class. Therefore, the inclusion Homeo_o $(D^2) \hookrightarrow$ Homeo_o (B^2) does not induce an isomorphism on bounded cohomology. It would be interesting to see if Homeo_o (D^3) has a non-trivial bounded cohomology.

To find a suitable resolution, we shall define semi-simplicial sets of fat chords:

Definition 5.4. A chord is an embedding of the closed arc D^1 into D^2 such that the endpoints ∂D^1 , but no other points of D^1 , are mapped to the circle ∂D^2 .

We fatten this definition by considering an orientation-preserving embedding $\varphi: D^1 \times \mathbf{R} \to D^2$ such that φ restricts to an embedding of $(\partial D^1) \times \mathbf{R}$ into ∂D^2 and such that $\varphi(B^1 \times \mathbf{R})$ lies in B^2 . Then let $[\varphi]$ denote the germ of φ around $D^1 \times \{0\}$, i.e. we say that φ and ψ have the same germ if the restrictions of φ and ψ to $D^1 \times (-\epsilon, \epsilon)$ are equal for some unspecified positive ϵ . We call the germ $[\varphi]$ a **fat chord** and denote the set of fat chords by FCh. The chord obtained by restricting φ to $D^1 \times \{0\}$ depends on $[\varphi]$ only and is called the **core** of $[\varphi]$. See Figure 1.



FIGURE 1. Two fat chords in D^2 with their cores.

Definition 5.5. We call two fat chords strictly transverse if

- their cores do intersect, and
- their cores are topologically transverse, and
- the endpoints of their cores are all distinct.

Consider a finite set of fat chords that are pairwise strictly transverse. Then the complement in B^2 of their cores is a finite union of open discs. The boundary of each such disc is partitioned into pieces of chords and pieces of the the original boundary $S^1 = \partial D^2$. The point of the first condition in Definition 5.5 is that there is at most one component from S^1 for each of these open discs in this partition. This restriction will later allow us to show that the stabilisers described in the next lemma are boundedly acyclic.

We write $\text{Homeo}(D^2; \text{near } D)$ for the subgroup of $\text{Homeo}(D^2)$ consisting of homeomorphisms that are trivial in some neighbourhood in D^2 of a boundary interval $D \subseteq \partial D^2$, that is, and embedded interval $D^1 \cong D \subseteq \partial D^2$. More precisely, this definition understands that an arbitrary such D has been chosen, but that each element of $\text{Homeo}(D^2; \text{near } D)$ can be trivial on a different neighbourhood of D in D^2 . The above definitions thus imply the following description.

Lemma 5.6. The stabiliser in Homeo_o (D^2) of any (n + 1)-tuple of pairwise strictly transverse fat chords is a finite product of groups, each isomorphic either

to Homeo $(D^2; \text{near } S^1)$ or to Homeo $(D^2; \text{near } D)$. Moreover, there are at most 2n+2 factors of the latter type.

On the other hand, given that chords may intersect a large number of times, there is no bound on the number of the factors of the Homeo $(D^2; \text{near } S^1)$ type.

Remark 5.7. The description recorded in Lemma 5.6 simplifies for the action of the subgroup Homeo(D^2 ; near S^1). In that case, the stabiliser of any tuple of pairwise strictly transverse fat chords is a finite power of Homeo(D^2 ; near S^1).

Next, we show that as the name suggests, strict transversality satisfies the condition of Definition 3.1.

Lemma 5.8. The relation of strict transversality is generic on FCh.

Proof. Given finitely many fat chords a_1, \ldots, a_k , the claim is that there is $a_0 \in FCh$ strictly transverse to a_i for all i > 0. We first outline the strategy of the proof, writing \dot{a}_i for the core of a_i .

There is no difficulty whatsoever in satisfying the first and last conditions of Definition 3.1. The substance of the lemma lies in ensuring the second condition, namely that the core \dot{a}_0 of the desired a_0 intersects transversely all given \dot{a}_i . Adjusting a_0 to be transverse to one a_i at a time can immediately be obtained by Schoenflies's theorem applied to \dot{a}_i , but the difficulty is that the various \dot{a}_i might intersect each other non-transversally, for instance in Cantor sets: we remind the reader that for i, j > 0, no transversality assumption is made for a_i relatively to a_j .

The strategy is to construct a_0 inductively, starting from a suitable fat chord a_0^0 satisfying the first and last conditions of Definition 3.1. We then use that the second condition is in fact "generic" in a topological sense. This allows us to perform a small perturbation (in the C^0 sense) of our first choice a_0^0 , to choose inductively a_0^i so that it is transverse to all $\{a_j\}_{j=1}^i$. We then define $a_0 = a_0^k$. This topological genericity of transversality for arcs is a standard fact, nicely explained in [Zac14]; we shall recall the details below.

The base a_0^0 of the induction must be chosen so that small C^0 perturbations do not make the intersections of cores empty. To that end, we choose a_0^0 in such a way that for each i > 0 some sub-arc of the core \dot{a}_0^0 intersects \dot{a}_i transversally. This can be done in steps as follows. Start from a boundary point distinct from all endpoints. By Schoenflies's theorem, \dot{a}_1 divides the disc into two discs and thus we can choose an initial (fat) arc from the boundary point to a point in the other disc component, crossing \dot{a}_1 once transversally and stopping at an interior point not lying on any \dot{a}_i . We then apply Schoenflies's theorem to the next core \dot{a}_2 and cross it transversally in the same fashion, repeating this argument until \dot{a}_k is crossed and ending on a new boundary point distinct from all endpoints. We need to avoid self-intersections of \dot{a}_0^0 , but this is possible since at every intermediate step the partially constructed arc \dot{a}_0^0 is retractable to the boundary.

Now we prepare the inductive argument. Given i, j > 0, consider the pairwise intersection $\dot{a}_i \cap \dot{a}_j$ of the cores and denote by K_{ij} the topological boundary $K_{ij} = \partial_{\dot{a}_i}(\dot{a}_i \cap \dot{a}_j)$ in \dot{a}_i of this intersection. Notice that $K_{ij} = K_{ji}$ holds because interior points of the intersection with respect to either \dot{a}_i or \dot{a}_j coincide. Consider the union $K_i = \bigcup_j K_{ij}$. Since the boundary of a closed set is always nowhere dense, K_i is a nowhere dense subset of the arc \dot{a}_i .

The inductive claim at step i (with $1 \le i \le k$) is that a_0^{i-1} admits an arbitrarily small perturbation a_0^i such that for all $1 \le j \le i$, the core \dot{a}_0^i is transverse to \dot{a}_j and avoids the set K_j .

The proof of this claim for i = 1 is a simpler version of the proof for i > 1, so we establish the latter while assuming that it already holds for all $1 \le j < i$. Consider

$$N = (\dot{a}_0^{i-1} \cap \dot{a}_i) \setminus \bigcup_{1 \le j < i} (\dot{a}_j \cap \dot{a}_i).$$

Since \dot{a}_0^{i-1} avoids $K_{ji} = K_{ij}$, we have in fact

$$N = (\dot{a}_0^{i-1} \cap \dot{a}_i) \smallsetminus \bigcup_{1 \le j < i} \operatorname{Int}_{a_i} (\dot{a}_j \cap \dot{a}_i)$$

so that N is a closed set and hence it has a neighbourhood V separating it from a_j for all $1 \leq j < i$. Now each sub-arc of \dot{a}_0^{i-1} in V can be perturbed within V to become transverse to \dot{a}_i using Schoenflies's theorem, and we can as well do it with the corresponding portion of the fat arc a_0^{i-1} . We can additionally ensure that the core of this new perturbation a_0^i avoids K_i since K_i is nowhere dense in the sub-arc of \dot{a}_i that we are crossing.

The reason that this completes the induction step is that outside N, the previous iteration \dot{a}_0^{i-1} was already transverse to \dot{a}_i because any intersection with \dot{a}_i in N would already occur in the interior of $\dot{a}_i \cap \dot{a}_j$ for some j < i. Now that the induction is complete, the final iteration a_0^k is the desired a_0 .

We consider now the semi-simplicial set FCh_{\bullet}^{\perp} of tuples of pairwise strictly transverse fat chords. Then Lemma 5.8 allows us to apply Proposition 3.2 and deduce:

Corollary 5.9. The semi-simplicial set FCh_{\bullet}^{\perp} is boundedly acyclic.

We shall also need to consider a somewhat technical subset of FCh and hence of FCh_{\bullet}^{\perp} , as follows.

Definition 5.10. A fat chord $[\varphi]$ is called **radial** (near the boundary) if we can choose $\varphi : D^1 \times \mathbf{R} \hookrightarrow D^2$ which is radial in a neighbourhood of $\{\pm 1\} \times \mathbf{R}$ in the following sense: there exists $\epsilon > 0$ such that

$$\varphi(\pm r, t) = r\varphi(\pm 1, t) \quad \forall \ 1 - \epsilon < r \le 1, \forall t.$$

We write $RFCh \subseteq FCh$ for the set of radial fat chords and $RFCh_{\bullet}^{\perp}$ for the corresponding sub-semi-simplicial set of FCh_{\bullet}^{\perp} .

This specific definition is somewhat arbitrary since we are working in the topological category; the point is only to choose some normalisation of the germ near the boundary. We note that that the arguments given for FCh can be repeated virtually unchanged to yield the following variation of Corollary 5.9.

Corollary 5.11. The semi-simplicial set $RFCh^{\perp}_{\bullet}$ is boundedly acyclic.

 \square

Remark 5.12. It is plausible to use more high powered transversality results in the topological settings ([KS77]) to prove an analogue of Corollary 5.9 and Corollary 5.11 in higher dimensions. We will, however, only be able to work in dimension two because the fact that the complementary regions in D^2 of simplices of FCh[⊥]_• is again a union of discs is, as we shall see, an important feature in the proof Theorem 5.1.

The relevance of the radial semi-simplicial subset hinges on the following observation:

Lemma 5.13. The inclusion map RFCh \rightarrow FCh descends to a bijection

RFCh /Homeo $(D^2; \text{near } S^1) \xrightarrow{\cong} \text{FCh} /\text{Homeo}(D^2; S^1).$

More generally, for all p it induces a bijection

 $\operatorname{RFCh}_p^{\perp}/\operatorname{Homeo}(D^2; \operatorname{near} S^1) \xrightarrow{\cong} \operatorname{FCh}_p^{\perp}/\operatorname{Homeo}(D^2; S^1).$

Proof. By the isotopy extension theorem for homeomorphisms in dimension 2 (see [Ham74, Thm. 1.7.10]) and the fact that $\text{Homeo}^{\tau}(D^2; S^1)$ as a topological group is contractible ([Ham74, Thm. 1.5.2]), we know that space of embeddings of chords with fixed ends in D^2 is contractible. In particular, if two fat chords have the same ends, they are isotopic relative to their ends. Hence, by an isotopy fixing the boundary pointwise, we can make the fat chords radial near the boundary and in fact we can make the support of this isotopy in any given neighbourhood of the boundary. Hence the above maps are surjective. For injectivity, note that if two radial fat chords have the same ends, they overlap near the boundary so they will be isotopic by an isotopy that is identity near the boundary.

In order to prove Theorem 5.1, we still need to deal with an auxiliary subgroup. We recall the notation Homeo $(D^2; \text{near } D)$ introduced before Lemma 5.6 for some boundary interval $D \subseteq S^1$.

Lemma 5.14. The group Homeo $(D^2; \text{near } D)$ satisfies the assumptions of Theorem 2.2 and hence is boundedly acyclic.



FIGURE 2. Z, Z_0 and the first displaced copy gZ_0 .

Proof of Lemma 5.14. It suffices to justify that we are in the situation of Theorem 2.2. To do so, we consider the following equivalent set-up (see Figure 5). Let Z be a square and let D be the union of three edges of this square. We call the remaining edge the **free edge** since elements of Homeo(D^2 ; near D) can restrict to non-trivial homeomorphisms of that edge. An important point is that each such edge homeomorphism will be supported in a compact subset of the interior of that edge.

Let now Z_0 be a smaller square inside Z whose free edge lies in the interior the free edge of Z. Then indeed we can find a "displacement" g and both conditions of Theorem 2.2 are satisfied.

Remark 5.15. In the first version of this article, we observed that Lemma 5.14 can be proved using the original Matsumoto–Morita method.

At this point, we can conclude from the description given in Lemma 5.6 that the stabiliser of any (n+1)-tuple of pairwise strictly transverse fat chords is a boundedly acyclic group since bounded acyclicity passes to products (Proposition 2.4). However, we shall need to know that the vanishing moduli of this bounded acyclicity does not depend on the combinatorics of chord intersections even though this combinatorics leads to unboundedly many factors in the product described in Lemma 5.6 even if we fix n. The desired uniformity is a consequence of Proposition 2.6, as follows.

Corollary 5.16. For every q > 0 there is a constant bounding the qth vanishing modulus of the stabiliser in Homeo_o(D^2) or in Homeo(D^2 ; near S^1) of any (n+1)-tuple of pairwise strictly transverse fat chords.

Proof. Let G be the product of countably many copies of $\text{Homeo}(D^2; \text{near } S^1) \cong$ Homeo_c(\mathbb{R}^2) and of Homeo($D^2; \text{near } D$). Both types of factors satisfy the assumptions of Theorem 2.2 and hence Lemma 2.3 implies that G is boundedly acyclic. According to Lemma 5.6 and Remark 5.7, all the stabilisers considered in the statement of Corollary 5.16 are subproducts of G. Therefore, the statement follows indeed from Proposition 2.6.

Proof of Theorem 5.1. For shorter notation, we write

 $G = \operatorname{Homeo}_{\circ}(D^2)$ and $G' = \operatorname{Homeo}(D^2; S^1)$

so that the restriction to S^1 yields an identification $G/G' = \text{Homeo}_{\diamond}(S^1)$.

We claim that the semi-simplicial set $\operatorname{FCh}^{\downarrow}_{\bullet}/G'$ is boundedly acyclic. In view of Lemma 5.13, it suffices to establish this for $\operatorname{RFCh}^{\downarrow}_{\bullet}/\operatorname{Homeo}(D^2; \operatorname{near} S^1)$ instead. We shall deduce this from Theorem 3.3. To that end, recall that $\operatorname{Homeo}(D^2; \operatorname{near} S^1) \cong \operatorname{Homeo}_c(\mathbb{R}^2)$ is boundedly acyclic by Matsumoto–Morita's theorem, while $\operatorname{RFCh}^{\downarrow}_{\bullet}$ is boundedly acyclic by Corollary 5.11. It thus remains to check the conditions on the stabilisers, which are granted by the Corollary 5.16. This confirms the claim.

The claim puts us in the position to apply Theorem 3.3 a second time, but to the action of G/G' on $\operatorname{FCh}^{\downarrow}_{\bullet}/G'$. The stabiliser of a (p+1)-tuple for this action corresponds to the stabiliser of 2(p+1) generic fat points in S^1 under the identification of G/G' with Homeo_o (S^1) . Thus, as in the Proof of Theorem 1.1, these stabilisers are boundedly acyclic (and there is only one isomorphism type when p is fixed). Therefore, Theorem 3.3 provides an isomorphism between the bounded cohomology of Homeo_o (S^1) and the bounded cohomology of the quotient of $\operatorname{FCh}^{\downarrow}_{\bullet}/G'$ by G/G', which is none other than $\operatorname{FCh}^{\downarrow}_{\bullet}/G$.

We invoke a third time Theorem 3.3, now for the G-action on $\operatorname{FCh}^{\downarrow}_{\bullet}$. That semisimplicial set is boundedly acyclic by Corollary 5.9. To justify the assumptions on the stabilisers, we recall the description of Lemma 5.6 and invoke again Corollary 5.16. We thus obtain an isomorphism between the bounded cohomology of Gand of $\operatorname{FCh}^{\downarrow}_{\bullet}/G$, which we previously identified with the bounded cohomology of Homeo_o(S^1).

It only remains to justify that this isomorphism is indeed induced by the restriction map $G \to \text{Homeo}_{\circ}(S^1)$, or equivalently by the quotient map $G \to G/G'$. The two isomorphisms produced by the second and third applications of Theorem 3.3 arise from parallel spectral sequences, connected by the morphism of spectral sequences induced by the quotient maps $\text{FCh}^{\perp}_{\bullet} \to \text{FCh}^{\perp}_{\bullet}/G'$ and $G \to G/G'$. Thus indeed the isomorphism is induced by restriction (which in particular preserves the ring structure of bounded cohomology); this completes the proof of Theorem 5.1.

Proof of Theorem 1.4. Our proof of Theorem 5.1 is set up in such a way that it can be followed equally well in the C^r case. We only need to replace our semi-simplicial sets $\mathrm{FCh}^{\perp}_{\bullet}$ and $\mathrm{RFCh}^{\perp}_{\bullet}$ by the corresponding sets of C^r germs. The main difference, viewed from the perspective of ordinary cohomology, is that the bounded acyclicity of stabilisers still holds but cannot be traced back to Matsumoto–Morita methods. Instead, we argue that Lemma 5.14 holds unchanged for (the connected component of) diffeomorphism groups because it relies on Theorem 2.2 and the latter does not discriminate according to regularity. The same holds for the bounded acyclicity of $\mathrm{Diff}^r_o(D^2; \mathrm{near} S^1)$.

6. Homeomorphisms and diffeomorphisms of the sphere S^n

Our goal in this section is to prove Theorem 1.8 which in particular solves Ghys's question (see Section 1.C) about invariants of flat S^3 -bundles. We first describe a strategy to determine the low degree bounded cohomology of $\text{Diff}^r_{\circ}(S^n)$ for $r \neq n+1$ and later we restrict to S^3 .

Before proceeding, we recall that a **quasimorphism** on a group G is a map $f: G \to \mathbf{R}$ such that the quantity |f(x) + f(x) - f(xy)| is bounded uniformly over $x, y \in G$. Every quasimorphism f lies at bounded distance from a unique **homo-geneous** quasimorphism \bar{f} , that is, a quasimorphism satisfying $\bar{f}(x^n) = n\bar{f}(x)$ for all $x \in G$ and $n \in \mathbf{Z}$, see e.g. [Bav91, §3.3]. A quasimorphism is called **trivial** if it is at bounded distance of a true homomorphism, or equivalently if \bar{f} is a true homomorphism. Noting that the quantity f(x) + f(x) - f(xy) is a coboundary (in the inhomogeneous model) and hence a bounded cocycle, one verifies readily that the space of quasimorphisms modulo trivial quasimorphisms is isomorphic to the kernel of the comparison map $H_b^2(G) \to H^2(G)$ from bounded to ordinary cohomology in degree two (compare again [Bav91, §3.3]).

In addition, Matsumoto–Morita have shown [MM85, Cor. 2.11] that this comparison map is injective if G is **uniformly perfect**, that is, if there is a bound Nsuch that every element of G can be expressed as a product of at most N commutators. We shall call this the "Matsumoto–Morita lemma" to distinguish it from their bounded acyclicity theorem. This lemma can also be checked directly on the above description in terms of quasimorphisms.

Back to S^n , Tsuboi [Tsu13, Tsu08] proved that Homeo_o (S^n) and Diff^r_o (S^n) for $r \neq n+1$ are uniformly perfect and hence Matsumoto–Morita's lemma implies that the map

$$H^2_b(\operatorname{Homeo}_{\circ}(S^n)) \longrightarrow H^2(\operatorname{Homeo}_{\circ}(S^n))$$

is injective and the same holds for $\text{Diff}^r_{\circ}(S^n)$ when $r \neq n+1$. By Thurston's theorem ([Thu74, Cor. (b) of Thm. 5]), we know that

$$H^2(\operatorname{Homeo}_{\circ}(S^n)) \cong H^2(\operatorname{BHomeo}_{\circ}^{\tau}(S^n)).$$

Given what is known of the right hand side, this already implies $H_b^2(\text{Homeo}_{\circ}(S^n)) = 0$ when n = 2, 3.

As announced in Theorem 1.10, we can establish this vanishing also for H_b^3 , for all n, and perhaps surprisingly for diffeomorphisms as well.

Proof of Theorem 1.10. We apply Theorem 3.3 to the action of the group $G = \text{Homeo}_{\circ}(S^n)$ or $\text{Diff}_{\circ}^r(S^n)$ on the semi-simplicial set X_{\bullet}^{\perp} of tuples of fat points in S^n with disjoint cores, exactly as for Theorem 1.1. We work with C^r fat point in the case of $\text{Diff}_{\circ}^r(S^n)$. The bounded acyclicity of X_{\bullet}^{\perp} is granted by Proposition 3.2. The difference with the case n = 1 is that we do not know the higher bounded cohomology of stabilisers and therefore we take N = 4 in Theorem 3.3. Thus, we need to establish the vanishing of $H_b^q(G_1)$ for all q > 0 with p + q < 4, where G_1 denotes the stabiliser of a point in X_p^{\perp} and $p \ge 0$. We note that condition (ii) of Theorem 3.3 then follows since there is only one type of stabiliser in X_p^{\perp} for each p.

The case p = 0 for homeomorphisms follows from the Matsumoto–Morita theorem since in that case $G_1 \cong \text{Homeo}_c(\mathbb{R}^n)$. In the C^r case we quote Theorem 1.6 instead.

Since H_b^1 vanishes for every group, the only remaining case is the vanishing of $H_b^2(G_1)$ when p = 1. Now $G_1 \cong \text{Homeo}_c(S^{n-1} \times \mathbf{R})$ or $\text{Diff}_c^r(S^{n-1} \times \mathbf{R})$ and we can apply Theorem 4.2.

Now Theorem 3.3 shows that the bounded cohomology of G coincides with that of the quotient semi-simplicial set X^{\perp}_{\bullet}/G . This quotient, however, is boundedly acyclic as soon as n > 1 because G acts transitively on tuples of fat points with distinct cores by a germ pasting argument as in the proof of Theorem 1.1. That is, given two (p + 1)-tuples of pairwise disjoint orientation-preserving embeddings of open balls into S^n , there is an orientation-preserving transformation of S^n in the corresponding regularity which sends each ball embedding among the first tuple to the corresponding embedding among the second tuple. The only difference with S^1 for this transitivity is that in dimension one the cyclic order of the points must be preserved.

Now we restrict to the case of S^3 and proceed to answer Ghys's question.

Proof of Theorem 1.8. The proof follows exactly the strategy of Theorem 1.10, but with N = 5 instead of N = 4, which requires the vanishing of H_b^2 of the stabiliser of a simplex in X_2 . But this stabiliser group is isomorphic to $\text{Diff}_c^r(P)$ where Pis obtained by removing 3 disjoint closed balls from S^3 . The mapping class group $\pi_0(\text{Diff}_c^r(P)^{\tau})$ is generated by the Dehn twists around sphere boundary components so it is a finite 2-torsion group (see [HM90, Lemma 3.2]). Hence, it is enough to show that $H_b^2(\text{Diff}_{c,o}^r(P)) = 0$. To do so, we first observe that the comparison map

(2)
$$H^2_b(\operatorname{Diff}^r_{c,\circ}(P)) \longrightarrow H^2(\operatorname{Diff}^r_{c,\circ}(P)),$$

is injective, and then we show that its image is trivial. By [FRY19, Thm. 4.1] we know that $\operatorname{Diff}_{c,\circ}^r(P)$ is uniformly perfect for $r \neq 4, 0$. Therefore, in these regularities, Matsumoto–Morita's lemma implies that the map (2) is injective. In particular, there is no nontrivial quasimorphism on $\operatorname{Diff}_{c,\circ}^r(P)$.

It is likely that $\text{Homeo}_{c,\circ}(P)$ is also uniformly perfect but we learned from Bowden that one can use automatic continuity of homogeneous quasimorphisms on diffeomorphism groups ([BHW21, Thm. A.5]) and the fact that homeomorphisms of 3-manifolds can be C^0 -approximated by diffeomorphisms ([Mun60, Thm. 6.3]) to deduce that if there were a nontrivial quasimorphisms on $\text{Homeo}_{c,\circ}(P)$, it would restrict to a nontrivial quasimorphisms on $\text{Diff}_{c,\circ}(P)$ which we know they do not exists by the uniform perfectness of $\text{Diff}_{c,\circ}(P)$. Hence, we also have the injectivity for the map

(3)
$$H^2_b(\operatorname{Homeo}_{c,\circ}(P)) \longrightarrow H^2(\operatorname{Homeo}_{c,\circ}(P)).$$

Now we need the following input about the cohomology of $\operatorname{Diff}_{c,\circ}^r(P)^{\tau}$.

Claim. $H^2(BDiff_{c,\circ}^r(P)^{\tau}) = 0.$

Proof of the claim: Since $\operatorname{Diff}_{c,\circ}^r(P)^{\tau}$ is a connected group, the classifying space $B\operatorname{Diff}_{c,\circ}^r(P)^{\tau}$ is simply connected. Hence, it is enough to show that $H_2(B\operatorname{Diff}_{c,\circ}^r(P)^{\tau};\mathbf{R}) = 0$. The action of $\operatorname{Diff}_{c,\circ}^r(S^2 \times \mathbf{R})^{\tau}$ on the space of embeddings $\operatorname{Emb}_{\circ}(D^3, S^2 \times \mathbf{R})$ gives rise to the fibration (see [Pal60])

$$\operatorname{Diff}_{c,\circ}^r(P)^{\tau} \to \operatorname{Diff}_{c,\circ}^r(S^2 \times \mathbf{R})^{\tau} \xrightarrow{\operatorname{res}} \operatorname{Emb}_{\circ}(D^3, S^2 \times \mathbf{R}),$$

where res is the restriction of a diffeomorphism to a fixed embedding of D^3 into $S^2 \times \mathbf{R}$. It is standard (e.g. [Kup19, Thm. 9.1.2]) to see that $\text{Emb}_{\circ}(D^3, S^2 \times \mathbf{R}) \simeq \text{Fr}^+(S^2 \times [0,1])$ where $\text{Fr}^+(-)$ means the oriented orthonormal frame bundle. Since the tangent bundle of $S^2 \times [0,1]$ is trivial (which is in fact true for all orientable 3-manifolds), we have $\text{Fr}^+(S^2 \times [0,1]) \cong S^2 \times [0,1] \times \text{SO}(3)$. Hence, if we deloop the above fibration, we obtain a fibration

(4)
$$S^2 \times [0,1] \times \mathrm{SO}(3)^{\tau} \to B\mathrm{Diff}_{c,\circ}^r(P)^{\tau} \to B\mathrm{Diff}_{c,\circ}^r(S^2 \times \mathbf{R})^{\tau}$$

Now, by Hatcher's theorem ([Hat83, Appendix]), we know that $B\text{Diff}_c(S^2 \times \mathbf{R})^{\tau} \simeq SO(3)^{\tau}$. Since $B\text{Diff}_{c,\circ}(S^2 \times \mathbf{R})^{\tau}$ is the universal cover for $B\text{Diff}_c(S^2 \times \mathbf{R})^{\tau}$, we conclude $B\text{Diff}_{c,\circ}(S^2 \times \mathbf{R})^{\tau} \simeq SU(2)$. Hence, to calculate $H_2(B\text{Diff}_{c,\circ}^{\tau}(P)^{\tau}; \mathbf{R})$, we

20

look at the second page of the Serre spectral sequence for the fibration (4), where we have a differential

$$d_2: E_{3,0}^2 = H_3(B\text{Diff}_{c,\circ}(S^2 \times \mathbf{R})^{\tau}; \mathbf{R}) \to E_{0,2}^2 = H_2(S^2 \times [0,1] \times \text{SO}(3)^{\tau}; \mathbf{R}),$$

whose cokernel is $H_2(BDiff_{c,\circ}^r(P)^{\tau}; \mathbf{R})$.

To determine this differential, note that $BDiff_{c,\circ}(S^2 \times \mathbf{R})^{\tau}$ is 2-connected, so we have $H_3(BDiff_{c,\circ}(S^2 \times \mathbf{R})^{\tau}; \mathbf{R}) = H_2(Diff_{c,\circ}(S^2 \times \mathbf{R})^{\tau}; \mathbf{R})$. Therefore, the differential d_2 is the map induced by the map

res:
$$\operatorname{Diff}_{c,\circ}^r (S^2 \times \mathbf{R})^\tau \to S^2 \times [0,1] \times \operatorname{SO}(3)^\tau$$

on the second homology.

To show that res induces a surjection on H_2 , first note that $H_2(S^2 \times [0,1] \times SO(3)^{\tau}; \mathbf{R})$ is generated by the S^2 factor. On the other hand, by Hatcher's theorem ([Hat83, Appendix]), we know that $\operatorname{Diff}_{c,o}^r(S^2 \times \mathbf{R})^{\tau} \simeq \Omega(SU(2)^{\tau})$ where $\Omega(SU(2)^{\tau})$ is the loop space on $SU(2)^{\tau}$. Hence, the map d_2 is induced by the natural map

$$\alpha: \Omega(\mathrm{SU}(2)^{\tau}) \to S^2 \times [0, 1],$$

which is given by the Hopf map to the slice $S^2 \times \{t\}$ at time t of the loop. It can be easily seen that α induces an isomorphism on H_2 . So d_2 is also an isomorphism. Therefore, we have $H_2(B\text{Diff}_{c,\circ}^r(P)^{\tau}; \mathbf{R}) = 0$.

We use this topological fact as an input to show that the image of the comparison map for $\operatorname{Diff}_{c,\circ}^r(P)$ is trivial. This is easier for $\operatorname{Homeo}_{c,\circ}(P)$ since by the classical theorem of Cerf ([Cer59]), the proof of Smale's conjecture ([Hat83, Appendix]) implies that $\operatorname{Homeo}_{c,\circ}(P)^{\tau} \simeq \operatorname{Diff}_{c,\circ}^r(P)^{\tau}$. Hence, by Thurston's theorem ([Thu74, Cor. (b) of Thm. 5]), we have

$$H^2(\operatorname{Homeo}_{c,\circ}(P)) = H^2(B\operatorname{Homeo}_{c,\circ}(P)^{\tau}) = 0.$$

Therefore, the injectivity of the comparison map (3) already implies that $H_b^2(\operatorname{Homeo}_{c,\circ}(P)) = 0$. For the case of $\operatorname{Diff}_{c,\circ}^r(P)$, we do not know whether $H^2(\operatorname{Diff}_{c,\circ}(P))$ vanishes so we need to work a little harder.

Claim. Let M^n be a n-manifold such that $H^2(BDiff_{c,\circ}^r(M)^{\tau}) = 0$ and suppose that we know that the comparison map (2) is injective for $Diff_{c,\circ}^r(M)$. Then for $r \neq n+1$ we have $H_b^2(Diff_{c,\circ}^r(M)) = 0$.

Proof of the claim: Recall that $BDiff_{c,o}^r(M)$ is homotopy fiber in the fibration

$$\overline{BDiff_{c,\circ}^r(M)} \to BDiff_{c,\circ}^r(M) \to BDiff_{c,\circ}^r(M)^{\tau}$$

Given that $BDiff_{c,\circ}^r(M)^{\tau}$ is simply connected, and we know that for $r \neq \dim(M) + 1$, we have $H^1(\overline{BDiff_{c,\circ}^r(M)}) = 0$ by [Mat84, Appendix] and also we have $H^2(BDiff_{c,\circ}^r(M)^{\tau}) = 0$ by the hypothesis, the spectral sequence for the above fibration implies that the map

$$H^2(\operatorname{Diff}_{c,\circ}^r(M)) \to H^2(\overline{B\operatorname{Diff}_{c,\circ}^r(M)}),$$

is injective. On the other hand, from Mather–Thurston's theorem for $r \neq \dim(M) + 1$, it follows that $H^2(\overline{BDiff}_{c,\circ}^r(M))$ only depends on dim(M) (see [Thu74, Second corollary at page 306] and the proof of the lemma in [Mat84, Appendix]). Therefore, the inclusion $\operatorname{Diff}_{c,\circ}^r(\mathbf{R}^n) \to \operatorname{Diff}_{c,\circ}^r(M)$ induces an isomorphism

$$H^2(\overline{B\mathrm{Diff}^r_{c,\circ}(M)}) \xrightarrow{\cong} H^2(\overline{B\mathrm{Diff}^r_{c,\circ}(\mathbf{R}^n)}).$$

It follows from the commutative diagram

that the left vertical map is also injective. Now the commutative diagram of comparison maps

implies that the left vertical map is also injective.

But we know that $H_b^2(\operatorname{Diff}_{c,\circ}^r(\mathbf{R}^n))$ is trivial by Theorem 4.2. Therefore, $H_b^2(\operatorname{Diff}_{c,\circ}^r(M))$ also vanishes.

Hence, by the claim, for $r \neq 4$, we obtain $H_b^2(\text{Diff}_{c,\circ}(P)) = 0$ which finishes the proof.

7. Further comments and questions

7.A. Homeomorphisms of certain geometric 3-manifolds. Burago–Ivanov– Polterovich ([BIP08, Section 3.3]) proved that $\text{Diff}_{\circ}(M)$ is uniformly perfect for any closed 3-manifold M. Therefore, as we have seen already by Matsumoto–Morita's lemma, the comparison map

$$H^2_b(\operatorname{Diff}_{\circ}(M)) \to H^2(\operatorname{Diff}_{\circ}(M))$$

is injective. Again by automatic continuity of homogeneous quasimorphisms on diffeomorphism groups ([BHW21, Thm. A.5]) and the fact that homeomorphisms of 3-manifolds can be C^{0} -approximated by diffeomorphisms ([Mun60, Thm. 6.3]), we can also conclude that

$$H^2_b(\operatorname{Homeo}_{\circ}(M)) \to H^2(\operatorname{Homeo}_{\circ}(M))$$

is injective. By Thurston's theorem ([Thu74, Cor. (b) of Thm. 5]), we know that

 $H^{2}(\operatorname{Homeo}_{\circ}(M)) = H^{2}(\operatorname{BHomeo}_{\circ}^{\tau}(M)).$

Thanks to the generalised Smale conjecture which has been extensively studied for many cases ([Hat83, Hat76, Iva76, Gab01, HKMR12, MS13]) and recently has been proved in the remaining cases by Bamler and Kleiner ([BK19, BK21]), we know the homotopy type of Homeo $^{\circ}_{\circ}(M)$ for a 3-manifold admitting a Thurston geometry. Combining a number of known facts about Homeo $^{\circ}(M)$, we prove:

Theorem 7.1. Let M be a closed hyperbolic 3-manifold or a closed Seifert fibered space whose fundamental group has \mathbf{Z} as its center.

Then $H_b^2(\text{Homeo}_{\circ}(M)) = 0.$

Proof. In the hyperbolic case, the generalised Smale conjecture was proved by Gabai ([Gab01]) which implies that $\operatorname{Homeo}_{\circ}^{\tau}(M) \simeq *$ as a topological group. Therefore, by the above argument we obtain $H_b^2(\operatorname{Homeo}_{\circ}(M)) = 0$. However for Seifert fibered manifold M, the generalised Smale conjecture implies that $\operatorname{Homeo}_{\circ}^{\tau}(M) \simeq (S^1)^k$ where k is the rank of the center of $\pi_1(M)$ except the case of the solid torus for which $\operatorname{Homeo}_{\circ}^{\tau}(M) \simeq S^1 \times S^1$ and the case of D^3 for which $\operatorname{Homeo}_{\circ}^{\tau}(D^3) \simeq \operatorname{SO}(3)^{\tau}$ (see the introduction of [MS13]). Hence, in our case of interest when

 $H^2(\operatorname{BHomeo}_{\circ}(M))$ is nontrivial it is isomorphic **R** which is generated by the Euler class induced by the circle action on the Seifert fibered space by rotating the circle fibers. K. Mann [Man20] showed that this Euler class is not a bounded class. Therefore, injectivity of $H^2_b(\operatorname{Homeo}_{\circ}(M)) \to H^2(\operatorname{Homeo}_{\circ}(M))$ implies that $H^2_b(\operatorname{Homeo}_{\circ}(M))$ is also zero in this case. \Box

7.B. Diffeomorphisms and homeomorphisms of \mathbb{R}^n . The algebraic properties of the automorphism groups of non-compact manifolds are more subtle to study. It would be interesting to apply the same techniques we used for spheres and the 2-disc to extract information about the bounded cohomology of diffeomorphisms and homeomorphisms of \mathbb{R}^n . The main motivation is to study which invariants of C^r -flat \mathbb{R}^n -bundles for $r \ge 0$ are bounded. For the C^0 -case, recall that BHomeo_o(\mathbb{R}^n)^{au} classifies oriented \mathbb{R}^n -microbundle and it has a nontrivial homotopy type (see [KRW20] for nontrivial characteristic classes of \mathbb{R}^n -microbundles). In particular, the Euler class $\mathscr{E} \in H^n(B$ Homeo_o(\mathbb{R}^n)^{au}) when n is even and all the Pontryagin classes \mathscr{P}_i for $i \le n/4$ are nontrivial. Hence, they also pull back nontrivially to $H^{\bullet}(\text{Homeo}_o(\mathbb{R}^n))$ since by McDuff's theorem ([McD80]) we know that

 $H^{\bullet}(B\mathrm{Homeo}_{\circ}(\mathbf{R}^n)^{\tau}) \cong H^{\bullet}(\mathrm{Homeo}_{\circ}(\mathbf{R}^n)).$

In fact, these classes are also nontrivial for C^r -flat \mathbb{R}^n -bundles for all regularities r > 0 by the following observation. Using a deep result of Segal ([Seg78, Prop. 1.3 and 3.1]), we know that there is a map

$$\operatorname{BDiff}^r_{\circ}(\mathbf{R}^n) \to \operatorname{B}\Gamma^r_{n,\circ},$$

which is a homology isomorphism, where $\mathrm{B}\Gamma_{n,\circ}^r$ is the classifying space of Haefliger structures for codimension *n* foliations that are transversely oriented, see Section 1 in [Seg78] for more details. On the other hand, there is map

$$\nu: \mathrm{B}\Gamma^r_{n,\circ} \to B\mathrm{GL}_n(\mathbf{R})^{\tau}_{\circ}$$

which classifies oriented normal bundles to the codimension n foliations. For all regularities, it is known that the map ν is at least (n+1)-connected, see Remark 1, Section II.6 in [Hae71]. Hence, in particular the induced map

$$H^{\bullet}(B\mathrm{GL}_n(\mathbf{R})^{\tau}_{\circ}) \to H^{\bullet}(\mathrm{Diff}^r_{\circ}(\mathbf{R}^n)),$$

is an isomorphism for $\bullet \leq n$. Therefore, the classes \mathscr{P}_i for $i \leq n/4$ are nontrivial and so is \mathscr{E} when n is even in $H^{\bullet}(\operatorname{Diff}^r_{\circ}(\mathbf{R}^n))$ for all r. However, for n = 2, Calegari ([Cal04, Thm. C]) showed that $\mathscr{E} \in H^2(\operatorname{Diff}^r_{\circ}(\mathbf{R}^2))$ is not a bounded class. We in fact show that there is no bounded invariant for flat \mathbf{R}^n -bundles over surfaces.

Theorem 7.2. We have $H_b^2(\text{Diff}_o^r(\mathbf{R}^n)) = 0$ for all n and all $r \neq n+1$.

Proof. As a consequence of Rybicki's theorem ([Ryb11, Thm. 1.2]) we know that $\operatorname{Diff}_{\circ}^{r}(\mathbf{R}^{n})$ is uniformly perfect for $r \neq n+1$. Therefore, Matsumoto–Morita's lemma ([MM85, Cor. 2.11]) implies that the map

$$H^2_b(\operatorname{Diff}^r_\circ(\mathbf{R}^n)) \longrightarrow H^2(\operatorname{Diff}^r_\circ(\mathbf{R}^n))$$

is injective for $r \neq n+1$. The remaining part of the proof consists in showing that the right hand side vanishes when $n \neq 2$, while for n = 2 it turns out that the non-zero elements of the right hand side are not in the image of the comparison map.

Recall the deep result of Segal ([Seg78, Prop. 1.3 and 3.1]) implies that there is a map $\operatorname{BDiff}_{\circ}^{r}(\mathbf{R}^{n}) \to \operatorname{B}\Gamma_{n,\circ}^{r}$ which is a homology isomorphism, where $\operatorname{B}\Gamma_{n,\circ}^{r}$ is the classifying space of Haefliger structures for codimension *n* foliations that are transversely oriented, see Section 1 in [Seg78] for more details. Since $r \neq n+1$, a theorem of Mather [Mat74, Section 7], implies that the natural map $\nu: \operatorname{B}\Gamma_{n,\circ}^{r} \to$ $\operatorname{BGL}_{n}^{r}(\mathbf{R})_{\circ}$ is at least (n+2)-connected. (Without the restriction $r \neq n+1$, we still know that the map ν is at least (n + 1)-connected, see Remark 1, Section II.6 in [Hae71].) Therefore, by combining Segal's theorem and Thurston's theorem, we have $H^2(\text{Diff}^r_{\circ}(\mathbf{R}^1)) = 0$ for $r \neq 2$ as desired for n = 1. For n = 2, we have $H^2(\text{Diff}^r_{\circ}(\mathbf{R}^2)) = \mathbf{R}$ generated by the Euler class for all r. On the other hand, a theorem of Calegari ([Cal04, Thm. C]) shows that the Euler class in $H^2(\text{Diff}^r_{\circ}(\mathbf{R}^2)) = \mathbf{R}$ is not a bounded class. Finally, for n > 2, since $H^2(\text{BGL}^\tau_n(\mathbf{R})_{\circ}; \mathbf{R}) = 0$, we have indeed $H^2(\text{Diff}^r_{\circ}(\mathbf{R}^n)) = 0$.

7.C. Questions. We shall end with few questions regarding the borderline of some of the cases we considered. For the case of spheres, it would be interesting to see if H_b^4 vanishes for S^n where $n \neq 1, 3$. In particular, the case of S^2 is already interesting. By Thurston's theorem ([Thu74, Cor. (b) of Thm. 5]) we know

 $H^4(\operatorname{Homeo}_{\circ}(S^2)) = H^4(\operatorname{BHomeo}_{\circ}^{\tau}(S^2)),$

and by Hamstrom's theorem ([Ham74]) we know that $\operatorname{Homeo}_{\circ}^{\tau}(S^2) \simeq \operatorname{SO}(3)^{\tau}$. Therefore, we have $H^4(\operatorname{Homeo}_{\circ}(S^2)) = \mathbb{R}$ generated by the Pontryagin class \mathscr{P}_1 in $H^4(\operatorname{BSO}(3)^{\tau})$.

Question 7.3. Is the first Pontryagin class bounded or even more generally is $H_b^4(\text{Homeo}_{\circ}(S^2))$ nontrivial?

Question 7.4. For n > 1, is there any nontrivial bounded class in $H^{\bullet}(\text{Homeo}_{\circ}(S^n))$?

In proving $\text{Homeo}_{\circ}(D^2) \to \text{Homeo}_{\circ}(S^1)$ we used a huge semi-simplicial set of transverse fat chords. There is an obvious way to generalize this semi-simplicial set to higher dimensions which is still boundedly acyclic. But then, given that the complement of simplices in D^n could be more complicated when n > 2, it is not clear whether the stabilisers are boundedly acyclic. One could try to define "smaller" semi-simplicial set whose stabilisers of its simplices are among the bounded acyclic groups. But it becomes harder to prove that the chosen semisimplicial set is boundedly acyclic. Hence, we pose the generalisation of our result about $\text{Homeo}_{\circ}(D^2)$ as a question.

Question 7.5. Does the restriction map $\text{Homeo}_{\circ}(D^n) \to \text{Homeo}_{\circ}(S^{n-1})$ induce an isomorphism on bounded cohomology for n > 2?

We generalised Calegari's theorem ([Cal04, Thm. C]) about the second bounded cohomology of $\operatorname{Diff}_{\circ}^{r}(\mathbf{R}^{2})$ by showing the vanishing $H_{b}^{2}(\operatorname{Diff}_{\circ}^{r}(\mathbf{R}^{n})) = 0$ for all n and all $r \neq n + 1$. So the higher degrees of bounded cohomology of $\operatorname{Diff}_{\circ}^{r}(\mathbf{R}^{2})$ remains to be determined.

Question 7.6. Is $\text{Diff}^r_{\circ}(\mathbf{R}^n)$ a boundedly acyclic group? Is $\mathscr{E} \in H^{2n}(\text{Diff}^r_{\circ}(\mathbf{R}^{2n}))$ a bounded class?

And finally, as we mentioned in the introduction, our proof of the unboundedness of the Euler class for oriented C^0 -flat S^3 -bundles is not constructive. It would be geometrically enlightening to find a constructive proof.

Question 7.7. Find explicit families of oriented C^0 -flat S^3 bundles over a given 4-manifold with unbounded Euler number.

References

- [Bav91] Christophe Bavard. Longueur stable des commutateurs. Enseign. Math. (2), 37(1-2):109–150, 1991.
- [BHW21] Jonathan Bowden, Sebastian Wolfgang Hensel, and Richard Webb. Quasi-morphisms on surface diffeomorphism groups. J. Amer. Math. Soc., 35(1):211–231, 2021.

- [BIP08] Dmitri Burago, Sergei Ivanov, and Leonid Polterovich. Conjugation-invariant norms on groups of geometric origin. In Groups of Diffeomorphisms: In honor of Shigeyuki Morita on the occasion of his 60th birthday, pages 221–250. Mathematical Society of Japan, 2008.
- [BK19] Richard H Bamler and Bruce Kleiner. Ricci flow and contractibility of spaces of metrics. arXiv preprint arXiv:1909.08710, 2019.
- [BK21] Richard H Bamler and Bruce Kleiner. Diffeomorphism groups of prime 3-manifolds. arXiv preprint arXiv:2108.03302, 2021.
- [BM12] Michelle Bucher and Nicolas Monod. The norm of the Euler class. Math. Ann., 353(2):523–544, 2012.
- [BM21] Michael Brandenbursky and Michał Marcinkowski. Bounded cohomology of transformation groups. Mathematische Annalen, pages 1–17, 2021.
- [Bot70] Raoul Bott. On a topological obstruction to integrability. In Proc. Symp. Pure Math, volume 16, pages 127–131, 1970.
- [Bot78] Raoul Bott. On some formulas for the characteristic classes of group-actions. In Differential topology, foliations and Gelfand-Fuks cohomology (Proc. Sympos., Pontificia Univ. Católica, Rio de Janeiro, 1976), volume 652 of Lecture Notes in Math., pages 25–61. Springer, Berlin, 1978.
- [Bow11] Jonathan Bowden. Flat structures on surface bundles. Algebr. Geom. Topol., 11(4):2207–2235, 2011.
- [Buc07] Michelle Bucher. Finiteness properties of characteristic classes of flat bundles. Enseign. Math. (2), 53(1-2):33-66, 2007.
- [Cal04] Danny Calegari. Circular groups, planar groups, and the Euler class. In Proceedings of the Casson Fest, volume 7 of Geom. Topol. Monogr., pages 431–491. Geom. Topol. Publ., Coventry, 2004.
- [Cer59] Jean Cerf. Groupes d'automorphismes et groupes de difféomorphismes des variétés compactes de dimension 3. Bull. Soc. Math. France, 87:319–329, 1959.
- [CFP96] James Welden Cannon, William J. Floyd, and Walter R. Parry. Introductory notes on Richard Thompson's groups. *Enseign. Math.* (2), 42(3-4):215–256, 1996.
- [Dup79] Johan L. Dupont. Bounds for characteristic numbers of flat bundles. In Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, 1978), pages 109–119. Springer, Berlin, 1979.
- [EHN81] David Eisenbud, Ulrich Hirsch, and Walter Neumann. Transverse foliations of Seifert bundles and self-homeomorphism of the circle. *Comment. Math. Helv.*, 56(4):638–660, 1981.
- [EP03] Michael Entov and Leonid Polterovich. Calabi quasimorphism and quantum homology. Int. Math. Res. Not., (30):1635–1676, 2003.
- [FFL21] Francesco Fournier-Facio and Yash Lodha. Second bounded cohomology of groups acting on 1-manifolds and applications to spectrum problems. Preprint, 2021.
- [FFLM21a] Francesco Fournier-Facio, Clara Löh, and Marco Moraschini. Bounded cohomology and binate groups. Preprint, 2021.
- [FFLM21b] Francesco Fournier-Facio, Clara Löh, and Marco Moraschini. Bounded cohomology of finitely generated groups: vanishing, non-vanishing, and computability. Preprint, 2021.
- [FRY19] Kazuhiko Fukui, Tomasz Rybicki, and Tatsuhiko Yagasaki. The uniform perfectness of diffeomorphism groups of open manifolds. arXiv preprint arXiv:1905.07664, 2019. [Gab01] David Gabai. The Smale conjecture for hyperbolic 3-manifolds: Isom $(M^3) \simeq$
- Diff (M^3) . Journal of Differential Geometry, 58(1):113–149, 2001.
- [GG04] Jean-Marc Gambaudo and Étienne Ghys. Commutators and diffeomorphisms of surfaces. Ergodic Theory Dynam. Systems, 24(5):1591–1617, 2004.
- [Ghy87] Étienne Ghys. Groupes d'homéomorphismes du cercle et cohomologie bornée. In The Lefschetz centennial conference, Part III (Mexico City, 1984), pages 81–106. Amer. Math. Soc., Providence, RI, 1987.
- [Ghy01] Étienne Ghys. Groups acting on the circle. Enseign. Math. (2), 47(3-4):329–407, 2001.
- [Gre87] Peter Greenberg. Classifying spaces for foliations with isolated singularities. Trans. Amer. Math. Soc., 304(1):417–429, 1987.
- [Gro82] Michaïl Gromov. Volume and bounded cohomology. Inst. Hautes Études Sci. Publ. Math., 56:5–99 (1983), 1982.
- [GS87] Étienne Ghys and Vlad Sergiescu. Sur un groupe remarquable de difféomorphismes du cercle. Comment. Math. Helv., 62(2):185–239, 1987.
- [Hae71] André Haefliger. Homotopy and integrability. In Manifolds-Amsterdam 1970, pages 133–163. Springer, 1971.

- [Hae79] A. Haefliger. Differential cohomology. In Differential topology (Varenna, 1976), pages 19–70. Liguori, Naples, 1979.
- [Hae10] André Haefliger. Differential cohomology. In Differential topology, pages 19–70. Springer, 2010.
- [Ham74] Mary-Elizabeth Hamstrom. Homotopy in homeomorphism spaces, TOP and PL. Bulletin of the American Mathematical Society, 80(2):207–230, 1974.
- $[{\rm Hat76}] \qquad {\rm Allen \ Hatcher. \ Homeomorphisms \ of \ sufficiently \ large \ P^2-irreducible \ 3-manifolds.} \\ Topology, 15(4):343-347, 1976.$
- [Hat83] Allen E. Hatcher. A proof of the Smale conjecture, $\text{Diff}(S^3) \simeq O(4)$. Ann. of Math. (2), 117(3):553–607, 1983.
- [HKMR12] Sungbok Hong, John Kalliongis, Darryl McCullough, and J. Hyam Rubinstein. Diffeomorphisms of elliptic 3-manifolds, volume 2055 of Lecture Notes in Mathematics. Springer, Heidelberg, 2012.
- [HM90] Allen Hatcher and Darryl McCullough. Finite presentation of 3-manifold mapping class groups. In Groups of Self-Equivalences and Related Topics, pages 48–57. Springer, 1990.
- [HT75] Morris W. Hirsch and William P. Thurston. Foliated bundles, invariant measures and flat manifolds. Ann. of Math. (2), 101:369–390, 1975.
- [IT82] Nikolai V. Ivanov and Vladimir G. Turaev. The canonical cocycle for the Euler class of a flat vector bundle. Dokl. Akad. Nauk SSSR, 265(3):521–524, 1982.
- [Iva76] N. V. Ivanov. Groups of diffeomorphisms of Waldhausen manifolds. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 66:172–176, 209, 1976. Studies in topology, II.
- [Jek12] Solomon Jekel. Powers of the Euler class. Adv. Math., 229(3):1949–1975, 2012.
- [Kot08] Dieter Kotschick. Stable length in stable groups. In Groups of diffeomorphisms, volume 52 of Adv. Stud. Pure Math., pages 401–413. Math. Soc. Japan, Tokyo, 2008.
- [KRW20] Alexander Kupers and Oscar Randal-Williams. On diffeomorphisms of evendimensional discs. arXiv preprint arXiv:2007.13884, 2020.
- [KS77] Robion C. Kirby and Laurence C. Siebenmann. Foundational essays on topological manifolds, smoothings, and triangulations. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1977. With notes by John Milnor and Michael Atiyah, Annals of Mathematics Studies, No. 88.
- [Kup19] Alexander Kupers. Lectures on diffeomorphism groups of manifolds. University Lecture Notes, 2019.
- [Lan94] Rémi Langevin. A list of questions about foliations. In Differential topology, foliations, and group actions (Rio de Janeiro, 1992), volume 161 of Contemp. Math., pages 59– 80. Amer. Math. Soc., Providence, RI, 1994.
- [Löh17] Clara Löh. A note on bounded-cohomological dimension of discrete groups. J. Math. Soc. Japan, 69(2):715–734, 2017.
- [Man20] Kathryn Mann. Unboundedness of some higher Euler classes. Algebraic & Geometric Topology, 20(3):1221–1234, 2020.
- [Mat71] John N. Mather. The vanishing of the homology of certain groups of homeomorphisms. *Topology*, 10:297–298, 1971.
- [Mat74] John N. Mather. Commutators of diffeomorphisms. Comment. Math. Helv., 49:512– 528, 1974.
- [Mat84] John N. Mather. A curious remark concerning the geometric transfer map. Comment. Math. Helv., 59(1):86–110, 1984.
- [Mat11] John N Mather. On the homology of Haefliger's classifying space. In Differential Topology, pages 71–116. Springer, 2011.
- [McD80] Dusa McDuff. The homology of some groups of diffeomorphisms. Commentarii Mathematici Helvetici, 55(1):97–129, 1980.
- [Mil58] John Milnor. On the existence of a connection with curvature zero. Comment. Math. Helv., 32:215–223, 1958.
- [Miz17] Tadayoshi Mizutani. On Thurston's construction of a surjective homomorphism $H_{2n+1}(B\Gamma_n, \mathbb{Z}) \to \mathbb{R}$. In Geometry, dynamics, and foliations 2013, volume 72 of Adv. Stud. Pure Math., pages 211–219. Math. Soc. Japan, Tokyo, 2017. Translated from the Japanese by Taro Asuke.
- [MM85] Shigenori Matsumoto and Shigeyuki Morita. Bounded cohomology of certain groups of homeomorphisms. Proceedings of the American Mathematical Society, 94(3):539– 544, 1985.
- [Mon01] Nicolas Monod. Continuous bounded cohomology of locally compact groups. Lecture Notes in Mathematics 1758, Springer, Berlin, 2001.

BOUNDED COHOMOLOGY OF HOMEOMORPHISM AND DIFFEOMORPHISM GROUPS 27

[Mon06] Nicolas Monod. An invitation to bounded cohomology. Proceedings of the international congress of mathematicians (ICM), Madrid, Spain, August 22-30, 2006. Volume II: Invited lectures. Zürich: European Mathematical Society, 1183–1211, 2006. [Mon13] Nicolas Monod. Groups of piecewise projective homeomorphisms. Proc. Natl. Acad. Sci. USA, 110(12):4524–4527, 2013. [Mon22] Nicolas Monod. Lamplighters and the bounded cohomology of Thompson's group. Geom. Funct. Anal., 32(3):662-675, 2022. [Mor84] Shigeyuki Morita. Nontriviality of the Gelfand-Fuchs characteristic classes for flat S¹-bundles. Osaka J. Math., 21:545–563, 1984. [Mor85] Shigeyuki Morita. Discontinuous invariants of foliations. In Foliations (Tokyo, 1983), volume 5 of Adv. Stud. Pure Math., pages 169–193. North-Holland, Amsterdam, 1985. [Mor17] Shigeyuki Morita. A few open problems on characteristic classes of foliations. In Geometry, dynamics, and foliations 2013, volume 72 of Adv. Stud. Pure Math., pages 221–237. Math. Soc. Japan, Tokyo, 2017. [MS04] Nicolas Monod and Yehuda Shalom. Cocycle superrigidity and bounded cohomology for negatively curved spaces. Journal of Differential Geometry, 67:395–455, 2004. [MS06] Nicolas Monod and Yehuda Shalom. Orbit equivalence rigidity and bounded cohomology. Ann. of Math. (2), 164(3):825-878, 2006. [MS13] Darryl McCullough and Teruhiko Soma. The Smale conjecture for Seifert fibered spaces with hyperbolic base orbifold. J. Differential Geom., 93(2):327-353, 2013. [Mun60] James Munkres. Obstructions to the smoothing of piecewise-differentiable homeomorphisms. Ann. of Math. (2), 72:521-554, 1960. [Nar16] Sam Nariman. On powers of the Euler class for flat circle bundles. Journal of Topology and Analysis, pages 1-6, 2016. [Nar20] Sam Nariman. A local to global argument on low dimensional manifolds. Transactions of the American Mathematical Society, 373(2):1307-1342, 2020. [Pal60] Richard S. Palais. Local triviality of the restriction map for embeddings. Comment. Math. Helv., 34:305-312, 1960. [Pol06] Leonid Polterovich. Floer homology, dynamics and groups. In Morse theoretic methods in nonlinear analysis and in symplectic topology, volume 217 of NATO Sci. Ser. II Math. Phys. Chem., pages 417-438. Springer, Dordrecht, 2006. [Ryb11] Tomasz Rybicki. Boundedness of certain automorphism groups of an open manifold. Geom. Dedicata, 151:175-186, 2011. [Seg78]Graeme Segal. Classifying spaces related to foliations. Topology, 17(4):367-382, 1978. [Smi] John Smillie. The Euler characteristic of flat bundles. unpublished manuscript. [Sul76] Dennis Sullivan. A generalization of Milnor's inequality concerning affine foliations and affine manifolds. Comment. Math. Helv., 51(2):183-189, 1976. William Thurston. Noncobordant foliations of S³. Bull. Amer. Math. Soc., 78:511-[Thu72] 514, 1972. [Thu74] William Thurston. Foliations and groups of diffeomorphisms. Bulletin of the American Mathematical Society, 80(2):304-307, 1974. Takashi Tsuboi. On the connectivity of the classifying spaces for foliations. In Alge-[Tsu89a] braic topology (Evanston, IL, 1988), volume 96 of Contemp. Math., pages 319-331. Amer. Math. Soc., Providence, RI, 1989. Takashi Tsuboi. On the foliated products of class C^1 . Ann. Math., 130:227–271, 1989. [Tsu89b] [Tsu00]Takashi Tsuboi. The Calabi invariant and the Euler class. Trans. Amer. Math. Soc., 352(2):515-524, 2000. [Tsu08] Takashi Tsuboi. On the uniform perfectness of diffeomorphism groups. In Groups of diffeomorphisms, volume 52 of Adv. Stud. Pure Math., pages 505-524. Math. Soc. Japan, Tokvo, 2008. Takashi Tsuboi. Homeomorphism groups of commutator width one. Proceedings of [Tsu13] the American Mathematical Society, 141(5):1839–1847, 2013. [Woo71] John W. Wood. Bundles with totally disconnected structure group. Comment. Math. Helv., 46:257-273, 1971. [Zac14]Zack. Topological transversality. Mathoverflow, July 16, 2014. École Polytechnique Fédérale de Lausanne (EPFL), CH–1015 Lausanne, Switzerland Email address: nicolas.monod@epfl.ch

Department of Mathematics, Purdue University, 150 N. University Street, West Lafayette, IN 47907-2067,

Email address: snariman@purdue.edu